

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1979

[Full Table of Content](#)



79/R/CS/133 Control of Pathogens - Maize

Rothamsted Research

Rothamsted Research (1980) *79/R/CS/133 Control of Pathogens - Maize* ; Yields Of The Field Experiments 1979, pp 129 - 130 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/R/CS/133

CONTROL OF PATHOGENS

Object: To study the effects of a range of chemicals on incidence of pathogens and yield of continuous maize - Long Hoos VI/VII 6.

Sponsors: A.J. Barnard, K.E. Fletcher, D.J. Hooper, D. Hornby, R.T. Plumb, T.D. Williams, J.C. Wilson.

The sixth year, forage maize.

For previous years see 74-78/R/CS/133.

Design: 3 randomised blocks of 9 plots split into 3.

Whole plot dimensions: 2.13 x 18.3.

Treatments: All combinations of:-

Whole plots

1. CHEMICAL	Chemicals applied annually except where stated:
NONE	None (2 plots per block)
ALDICARB	Aldicarb, 4.5 kg as granules to seedbed
BENOMYL	Benomyl, 11.2 kg as dust to seedbed
DAZOMET	Dazomet, 450 kg as granules in early spring (not applied 1975 & 1979)
PERMETH	Permethrin, 0.15 kg as foliar spray on 3 Aug (1979 only)
PHORATE	Phorate, 1.68 kg as granules to seedbed
PIRIMICA	Pirimicarb, 0.14 kg as foliar spray on 3 Aug (1979 only)
BE/DA/PH	Benomyl + dazomet (not applied 1975 & 1979) + phorate, at above rates and times

Sub plots

2. N Nitrogen fertiliser (kg N):

50
100
150

NOTE: Treatment sprays were applied in 340 l.

Basal applications: Weedkiller: Atrazine at 1.7 kg in 340 l.

Seed: Caldera 535, sown at 100,000 seeds per hectare.

Cultivations, etc.: - Ploughed: 9 Feb, 1979. Spring-tine cultivated: 16 May. Seedbed treatments applied, power harrowed, seed sown: 8 June. Weedkiller applied: 11 June. Harvested by hand: 8 Nov.

NOTES: (1) Germination was assessed in June.
(2) Frit fly (*Oscinella frit*) damage was assessed.
(3) N percentages in harvest produce were determined.

79/R/CS/133

FORAGE MAIZE TONNES/HECTARE

***** TABLES OF MEANS *****

	N	50	100	150	MEAN
CHEMICAL					
NONE		5.73	5.94	5.87	5.85
ALDICARB		6.67	6.59	8.14	7.13
BENQMYL		6.08	6.38	6.87	6.44
DAZOMET		5.90	7.08	6.84	6.61
PERMETH		5.83	5.97	6.50	6.10
PHORATE		6.02	6.28	6.17	6.16
PIRIMICA		6.60	6.57	7.25	6.81
BE/DA/PH		6.74	6.86	7.36	6.99
MEAN		6.14	6.40	6.76	6.44

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHEMICAL	N	CHEMICAL	
			N	
SED	0.370		0.528	MIN REP
	0.321	0.153	0.457	MAX-MIN
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:				
CHEMICAL			0.460	MIN REP
			0.326	MAX REP

CHEMICAL
 MAX REP WITHIN NONE
 MAX-MIN NONE V REMAINDER
 MIN REP REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	17	0.454	7.0
BLOCK.WP.SP	38	0.564	8.8

GRAIN MEAN DM% 20.6

SUB PLOT AREA HARVESTED 0.00059