Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1979



Full Table of Content

# 79/R/CS/211 Factors Affecting Eyespot - Wheat

# **Rothamsted Research**

Rothamsted Research (1980) 79/R/CS/211 Factors Affecting Eyespot - Wheat; Yields Of The Field Experiments 1979, pp 219 - 221 - DOI: https://doi.org/10.23637/ERADOC-1-45

#### 79/R/CS/211

# FACTORS AFFECTING EYESPOT

Object: To study the effects of a range of treatments on the incidence of eyespot (Pseudocercosporella herpotrichoides) and on the yield of wheat - Meadow.

Sponsors: R.D. Prew, A. Bainbridge.

The second year, wheat.

For previous year see 78/R/CS/211.

Design: 2 randomised blocks of 2 whole plots split into 6 sub plots split into 3 sub sub plots.

Whole plot dimensions: 94.0 x 9.14.

Treatments: All combinations of:-

Whole plots

1. STRAW Treatment of straw in autumn 1977 & 1978:

BURNT Burnt on site after spreading

CARTED Baled and carted off

Sub plots

2. DRILLING Cultivations and drilling in autumn 1977 & 1978:

CNVNTIAL Cultivated and conventionally drilled

DIRECT Uncultivated, direct drilled

3. SOW DATE Dates of sowing, autumn 1978:

26 SEPT 26 September 12 OCT 12 October 10 NOV 10 November

Sub sub plots

4. SEEDRATE Seed rates, autumn 1978 (kg):

100

150

200

NOTE: All treatments involving the combinations of STRAW CARTED & DRILLING DIRECT were severely infested with sterile brome grass and were cut green, no yields, on 18 June. Regrowth was cut again on 23 July.

#### 79/R/CS/211

Basal applications: Manures: (10:23:23) at 250 kg, combine drilled. 'Nitro-Chalk' at 500 kg. Weedkillers: Paraquat at 0.56 kg ion in 220 l. Methabenzthiazuron at 1.6 kg in 220 l. Mecoprop at 2.5 kg in 220 l.

Seed: Kador.

Oultivations, etc.:- Straw treatments applied: 9-12 Sept, 1978. Unburnt stubble cut and removed: 20 Sept. Paraquat applied: 22 Sept. CNVNTIAL treatments ploughed, and CNVNTIAL 26 SEPT treatments rotary harrowed, seed sown for all 26 SEPT treatments, CNVNTIAL treatments tine harrowed in, DIRECT treatments disc harrowed in and rolled: 26 Sept. CNVNTIAL 12 OCT treatments rotary harrowed, all 12 OCT treatments sown, CNVNTIAL treatments tine harrowed in, DIRECT treatments disc harrowed in and both treatments rolled: 12 Oct. CNVNTIAL 10 NOV treatments rotary harrowed, all 10 NOV treatments sown, CNVNTIAL treatments tine harrowed in, DIRECT treatments disc harrowed in and both treatments rolled: 10 Nov. Methabenzthiazuron applied: 11 Nov. N applied: 18 Apr, 1979. Mecoprop applied: 9 May. Combine harvested: 30 Aug. Previous crops: Wheat 1977 and 1978.

NOTE: Plants were assessed for infection with eyespot and the incidence of eyespot spores was measured throughout the year. Take-all was assessed at harvest.

#### GRAIN TONNES/HECTARE

#### \*\*\*\* TABLES OF MEANS \*\*\*\*

STRAW	SCW DATE DRILLING	26 SEPT	12 OCT	10 NOV	MEAN
BUR NT BUR NT	CNVNTIAL DIRECT CNVNTIAL	5.14 5.96 5.94	5.29 5.75 5.44	4.53 5.58 4.90	4.98 5.76 5.43
	MEAN	5.68	5.49	5.00	5.39
STRAW	SEEDRATE	100	150	200	MEAN
BUR NT BUR NT	CNVNTIAL DIRECT CNVNTIAL	4.90 5.59 5.36	4.99 5.68 5.48	5.07 6.01 5.44	4.98 5.76 5.43
	MEAN	5.28	5.38	5.51	5.39
SEEDRATE	100	150	200	MEAN	
SOW DATE 26 SEPT 12 OCT 10 NOV	5.73 5.30 4.81	5.52 5.57 5.05	5.78 5.60 5.14	5.68 5.49 5.00	
MEAN	5.28	5.38	5.51	5.39	

# 79/R/CS/211

# GRAIN TONNES/HECTARE

### \*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

STRAW	DRILLING	SEEDRATE SOW DATE	100	150	200
BURNT	CNVNTIAL	26 SEPT	5.29	4.99	5.14
		12 OCT	5.18	5.44	5.24
		10 NOV	4.23	4.55	4.82
BURNT	DIRECT	26 SEPT	5.72	5.59	6.55
		12 OCT	5.61	5.79	5.85
		10 NOV	5.45	5.65	5.63
CARTED	CNVNTIAL	26 SEPT	6.18	5.99	5.65
		12 OCT	5.13	5.47	5.72
		10 NOV	4.77	4.97	4.96

# \*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	SOW DATE	SEEDRATE	STRAW* DRILLING	SOW DATE SEEDRATE
SED	0.255	0.127	0.255	0.312
	COMPARING MEANS V			0.312
SOW DATE				0.221

TABLE	STRAW* DRILLING SOW DATE	STRAW* DRILLING SEEDRATE	STRAW* DRILLLING SEEDRATE SOW DATE
SED	0.442	0.312	0.541
EXCEPT WHEN STRAW.DRI	COMPARING MEANS	WITH SAME LEV 0.221	VEL(S) OF:
STRAW.DRI	LLING.SOW DATE		0.382

<sup>\*</sup> WITHIN THE SAME LEVEL OF STRAW ONLY

# \*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP.SP	7	0.442	8.2
BLOCK.WP.SP.SSP	18	0.382	7.1

GRAIN MEAN DM% 87.7

SUB PLOT AREA HARVESTED

DRILLING CNVNTIAL 0.00253
DRILLING DIRECT 0.00248