

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1979

[Full Table of Content](#)



### 79/R/CS/241 Pesticides and Slot-seeding - Ryegrass

#### Rothamsted Research

Rothamsted Research (1980) *79/R/CS/241 Pesticides and Slot-seeding - Ryegrass* ; Yields Of The Field Experiments 1979, pp 236 - 240 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/R/CS/241

PESTICIDES AND SLOT-SEEDING

Object: To study the effects of aldicarb and phorate, applied at sowing, on the incidence of pests and on the survival and yield of ryegrass sown into an existing grass sward by a slot-seeding drill - Little Knott II.

Sponsors: I.F. Henderson, A.M. Spaul, R.W. Gibson.

The first year, grass.

Design: 3 randomised blocks of 7 plots split into 3.

Whole plot dimensions: 2.0 x 24.0.

Treatments: All combinations of:-

Whole plots

1. TREATMNT	Treatments, applied to old grass:
0 0 0	None
0 S 0	Slot seeder used not sowing seed
G S R	Old grass killed with glyphosate, RvP ryegrass sown by slot-seeder
0 S B	Baroldi ryegrass sown by slot-seeder
0 S M	Grasslands Manawa ryegrass sown by slot-seeder
0 S R	RvP ryegrass sown by slot-seeder
0 S S	S.24 ryegrass sown by slot-seeder

Sub plots

2. PESTICIDE	Pesticides applied at sowing:-
NONE	None
ALDICARB	Aldicarb at 10 kg
PHORATE	Phorate at 10 kg

- NOTES: (1) The slot-seeder cut slots in the sward 2.5 cm deep and 30 cm apart, using knife coulters. Seed was sown at 12 kg.  
(2) Glyphosate to G S R was applied at 3.0 kg in 1000 l on 4 Apr, 1979.  
(3) One replicate of each of the treatment combinations 0 S B ALDICARB, 0 S B NONE, 0 0 0 NONE, 0 0 0 ALDICARB, was sprayed with glyphosate in error. Estimated values were used in the analysis.

Basal applications: Manures: (20:14:14) at 125 kg combine drilled. (Applied by hand in rows 30 cm apart to 0 0 0). (25:0:16) at 200 kg after each cut except the last.

Cultivations, etc.:- Seed sown: 6 Apr, 1979. Cut: 30 May, 27 June, 25 July, 22 Aug, 18 Sept, 17 Oct. NK applied: 6 June, 28 June, 26 July, 24 Aug, 19 Sept.

NOTE: Samples of harvested produce were sorted and weighed to determine amounts of original sward, dicotyledonous weeds and sown species.

79/R/CS/241

1ST CUT (30/5/79) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PESTCIDE TREATMNT	NONE	ALDICARB	PHORATE	MEAN
O O O	3.08	3.10	3.44	3.21
O S O	2.84	2.39	2.74	2.65
G S R	0.32	0.24	0.44	0.33
O S B	2.45	2.98	2.63	2.69
O S M	2.65	2.95	2.73	2.78
O S R	2.23	2.56	3.10	2.63
O S S	2.56	2.54	2.50	2.53
MEAN	2.31	2.39	2.51	2.40

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT	PESTCIDE	TREATMNT PESTCIDE
SED	0.170	0.092	0.261
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
TREATMNT			0.243

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.208	8.7
BLOCK.WP.SP	24	0.298	12.4

2ND CUT (27/6/79) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PESTCIDE TREATMNT	NONE	ALDICARB	PHORATE	MEAN
O O O	1.35	1.33	1.23	1.30
O S O	1.69	1.41	1.76	1.62
G S R	1.63	1.90	2.08	1.87
O S B	1.41	1.38	1.79	1.52
O S M	1.38	1.50	1.45	1.44
O S R	1.53	1.68	1.49	1.56
O S S	1.54	1.52	1.69	1.58
MEAN	1.50	1.53	1.64	1.56

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT	PESTCIDE	TREATMNT PESTCIDE
SED	0.204	0.062	0.244
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
TREATMNT			0.165

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.250	16.0
BLOCK.WP.SP	24	0.202	13.0

79/R/CS/241

3RD CUT (25/7/79) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PESTCIDE TREATMNT	NONE	ALDICARB	PHORATE	MEAN
0 0 0	1.09	0.76	0.86	0.90
0 S 0	0.81	0.82	0.92	0.85
G S R	1.20	1.43	1.36	1.33
0 S B	0.84	0.96	0.75	0.85
0 S M	0.57	0.72	0.75	0.68
0 S R	0.73	0.81	0.89	0.81
0 S S	0.59	0.91	0.87	0.79
MEAN	0.83	0.91	0.91	0.89

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT	PESTCIDE	TREATMNT PESTCIDE
SED	0.109	0.046	0.148
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
TREATMNT			0.123

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.133	15.0
BLOCK.WP.SP	24	0.150	17.0

4TH CUT (22/8/79) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PESTCIDE TREATMNT	NONE	ALDICARB	PHORATE	MEAN
0 0 0	1.14	1.10	1.13	1.13
0 S 0	0.90	1.16	1.26	1.11
G S R	1.35	1.50	1.53	1.46
0 S B	1.18	1.15	1.27	1.20
0 S M	1.05	1.20	1.12	1.12
0 S R	1.15	1.27	1.22	1.21
0 S S	0.84	1.17	1.33	1.11
MEAN	1.09	1.22	1.27	1.19

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT	PESTCIDE	TREATMNT PESTCIDE
SED	0.075	0.058	0.147
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
TREATMNT			0.155

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.092	7.8
BLOCK.WP.SP	24	0.189	15.9

79/R/CS/241

5TH CUT (18/9/79) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PESTCIDE TREATMNT	NONE	ALDICARB	PHORATE	MEAN
O O O	0.79	0.56	0.68	0.68
O S O	0.72	0.80	0.69	0.74
G S R	0.68	0.69	0.76	0.71
O S B	0.80	0.79	0.64	0.74
O S M	0.60	0.68	0.69	0.66
O S R	0.71	0.82	0.86	0.80
O S S	0.49	0.70	0.72	0.64
MEAN	0.68	0.72	0.72	0.71

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT	PESTCIDE	TREATMNT PESTCIDE
SED	0.077	0.034	0.106
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF: TREATMNT			0.089

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.094	13.3
BLOCK.WP.SP	24	0.109	15.4

6TH CUT (17/10/79) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PESTCIDE TREATMNT	NONE	ALDICARB	PHORATE	MEAN
O O O	0.47	0.72	0.62	0.60
O S O	0.54	0.46	0.53	0.51
G S R	0.53	0.49	0.55	0.52
O S B	0.40	0.41	0.56	0.46
O S M	0.57	0.54	0.56	0.56
O S R	0.44	0.53	0.57	0.52
O S S	0.50	0.51	0.57	0.52
MEAN	0.49	0.52	0.57	0.53

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT	PESTCIDE	TREATMNT PESTCIDE
SED	0.041	0.029	0.074
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF: TREATMNT			0.076

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.050	9.6
BLOCK.WP.SP	24	0.093	17.6

79/R/CS/241

TOTAL OF 6 CUTS DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PESTCIDE TREATMNT	NONE	ALDICARB	PHORATE	MEAN
O O O	7.93	7.57	7.96	7.82
O S O	7.50	7.03	7.89	7.48
G S R	5.71	6.24	6.72	6.23
O S B	7.08	7.66	7.65	7.46
O S M	6.82	7.58	7.30	7.23
O S R	6.78	7.68	8.14	7.53
O S S	6.51	7.35	7.66	7.18
MEAN	6.90	7.30	7.62	7.28

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT	PESTCIDE	TREATMNT PESTCIDE
SED	0.222	0.143	0.381
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
TREATMNT			0.380

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.272	3.7
BLOCK.WP.SP	24	0.465	6.4

1ST CUT MEAN DM% 17.2  
 2ND CUT MEAN DM% 17.8  
 3RD CUT MEAN DM% 26.2  
 4TH CUT MEAN DM% 18.8  
 5TH CUT MEAN DM% 23.1  
 6TH CUT MEAN DM% 18.0  
 TOTAL OF 6 CUTS MEAN DM% 20.2

AVERAGE PLOT AREA HARVESTED 0.00071