Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1979



Full Table of Content

## 79/R/B/5 N and Growth Regulator - Barley

### **Rothamsted Research**

Rothamsted Research (1980) 79/R/B/5 N and Growth Regulator - Barley; Yields Of The Field Experiments 1979, pp 295 - 296 - **DOI:** https://doi.org/10.23637/ERADOC-1-45

#### 79/R/B/5

#### WINTER BARLEY

#### N & GROWTH REGULATOR

Object: To study the effects of a growth regulator and rates and times of applying nitrogen on the yield of winter barley - Long Hoos I/II.

Sponsors: F.V. Widdowson, J.F. Jenkyn.

Design: 4 randomised blocks of 13 plots.

Whole plot dimensions: 2.13 x 9.14.

Treatments: All combinations of:-

1. E N TIME Times of applying early nitrogen:

FEB/MAR 30 kg of total early N applied 1 Mar, 1979, remainder 2 Apr.

MAR All early N applied 2 Apr.

2. E N RATE Total early nitrogen rate (kg N):

60 90

3. L N G Late nitrogen (kg N) and growth regulator:

NONE None

30 APR 30 kg applied 25 Apr. No growth regulator

30 APR+G 30 kg applied 25 Apr. Mepiquat chloride + ethephon (as 'Terpal' applied at 2.5 1) in 280 1

plus one extra plot:

90F/M+G 90 kg N total applied: 30 kg N 1 Mar, 60 kg N 2 Apr.

Mepiquat chloride + ethephon applied at above rate
on 25 May

NOTES: (1) Planned dates of applying early nitrogen treatments were not achieved because of wet weather.

(2) The guard areas between sides of each plot were sown to winter barley, variety Hoppel, and used for the experiment 'Sowing Dates, Mildew Control and Growth Study' (see 79/R/B/1).

Basal applications: Manures: (0:20:20) at 310 kg. Weedkillers: Mecoprop at 2.5 l in 220 l. Irrigation: 25 mm water.

Seed: Athene, sown at 160 kg.

Cultivations, etc.:- Ploughed: 22 Sept, 1978. Rolled: 27 Sept. PK applied, rotary harrowed: 4 Oct. Seed sown: 6 Oct. Irrigated: 9 Nov. Weedkiller applied: 9 May, 1979. Harvested: 6 Aug. Previous crops: Spring barley 1977, Winter beans 1978.

NOTE: Soil samples were taken in spring to a depth of 90 cm to determine mineral N content. Nitrogen percentages of grain were measured. Leaf disease and crop height were assessed in late June.

79/R/B/5					
GRAIN	TONNES	/HE	CTARE		
****	TABLES	OF	MEANS	****	

INDUID	OI IIDIIIIO		
E N RATI		90	MEAN
E N TIMI FEB/MAI	R 8.30	8.67	8.49
MAI	R 8.46	8.98	8.72
MEAT	0 20	0 00	9 60

	FEB/MAR MAR	8.46	8.98		8.49	
	MEAN	8.38	8.82		8.60	
	L N G E N TIME	NONE	30 APR	30	APR+G	MEAN
	FEB/MAR MAR	7.90 8.44	8.63 8.65		8.93 9.05	8.49 8.72
	MEAN	8.17	8.64		8.99	8.60
	L N G E N RATE	NONE	30 APR	30	APR+G	MEAN
	60 90	7.77 8.57	8.44 8.84		8.93 9.06	8.38 8.82
	MEAN	8.17	8.64		8.99	8.60
	E N TIME	L N G E N RATE	NC	NE	30 APR	30 APR+G
	FEB/MAR	60 90		58 23	8.45 8.80	
	MAR	60		97	8.42	

90F/M+G 8.62

GRAND MEAN 8.60

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

90

TABLE	E N TIME	E N RATE	L N G	E N TIME E N RATE
SED	0.119	0.119	0.146	0.168
TABLE	E N TIME L N G	E N RATE L N G	E N TIME E N RATE L N G & 90F/M+G	

8.92

8.88

SED 0.206 0.206 0.291

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM DF SE CV% BLOCK.WP 36 0.412 4.8

GRAIN MEAN DM% 86.3

PLOT AREA HARVESTED 0.00195