Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1979



Full Table of Content

79/R/B/8 and 79/W/B/8 Pythium Control - Barley

Rothamsted Research

Rothamsted Research (1980) 79/R/B/8 and 79/W/B/8 Pythium Control - Barley; Yields Of The Field Experiments 1979, pp 302 - 304 - DOI: https://doi.org/10.23637/ERADOC-1-45

79/R/B/8 and 79/W/B/8

SPRING BARLEY

PYTHIUM CONTROL

Object: To study the effects of two fungicides and two methods of application on the incidence of Pythium on roots and on the yield of spring barley - Rothamsted (R) Gt. Harpenden I and Woburn (W) Stackyard C.

Sponsor: G.A. Salt.

Design: 3 randomised blocks of 6 plots.

Whole plot dimensions: 4.27 x 9.14.

Treatments: All combinations of:-

FUNGCIDE Fungicides:

ALIETTE 'Aliette' (Aluminium tris (ethyl phosphonate))
'CGA 'CGA 48988' (DL-methyl N-(2,6 dimethyl phenyl)-N(2-methoxyacetyl)alaninate

2. FUNGMETH Method of applying fungicides:

FOLIAR Foliar spray

SEEDBED Rotavated into the seedbed

plus one extra treatment, duplicated:

NONE

NOTES: (1) 'Aliette' was applied at 15 kg in the seedbed and at 0.68 kg as a foliar spray.

(2) 'CGA 48988' was applied at 1.5 kg in the seedbed and at 0.07 kg as a foliar spray.

Basal applications:

Gt. Harpenden I (R): Manures: (25:0:16) at 450 kg, combine drilled. Weedkillers: Bromoxynil with ioxynil ('Oxytril CM' at 1.4 kg) and mecoprop at 1.7 kg in 220 l.

Stackyard C (W): Manures: (25:0:16) at 450 kg. Weedkillers: Dicamba with mecoprop and MCPA ('Banlene Plus' at 4.9 kg in 250 1).

Seed: Porthos, dressed with ethirimol, sown at 160 kg at both sites.

Cultivations, etc .:-

Gt. Harpenden I (R): Ploughed: 3 Aug, 1978. Deep-tine cultivated: 10 Aug. Heavy spring-tine cultivated twice: 24 Aug, 7 Sept. Spring-tine cultivated: 30 Apr, 1979. Seedbed treatments applied, spike rotary cultivated all plots, seed sown: 3 May. Weedkillers applied: 4 June. Foliar treatments applied: 12 June. Combine harvested: 5 Sept. Previous crops: Ryegrass 1977, 1978.

79/R/B/8 and 79/W/B/8

Stackyard C (W): Ploughed: 22 Nov, 1978. Spring-tine cultivated with crumbler attached, three times: 17 Apr, 1979, 30 Apr, 8 May. NK applied: 27 Apr. Seedbed treatments applied, rotary cultivated, all plots sown: 8 May. Weedkillers applied: 6 June. Foliar treatments applied: 18 June. Combine harvested: 6 Sept. Previous crops: Fallow 1977, 1978.

NOTE: Crop samples were taken in July for assessment of infection by Pythium spp. and other soilborne fungi.

79/R/B/8 GT. HARPENDEN I(R)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE FUNGMETH	ALIETTE	CGA	MEAN
FOLIAR SEEDBED	4.39 4.53	4.15 4.46	4.27
MEAN	4.46	4.31	4.38

NONE 4.48

GRAND MEAN 4.42

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGCIDE	FUNGMETH	FUNGCIDE FUNGMETH & NONE
SED	0.213	0.213	0.302 0.261*

^{*} FOR COMPARISONS INVOLVING NONE ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	11	0.370	8.4

GRAIN MEAN DM% 86.0

PLOT AREA HARVESTED 0.00195

79/W/B/8 STACKYARD C(W)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE	ALIETTE	CGA	MEAN
FUNGMETH			
FOLIAR	2.20	2.41	2.31
SEEDBED	2.51	2.37	2.44
MEAN	2.36	2.39	2.37

NONE 2.59

GRAND MEAN 2.45

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGCIDE	FUNGMETH	FUNGCIDE FUNGMETH & NONE
SED	0.280	0.280	0.396

^{*} FOR COMPARISONS INVOLVING NONE ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	6	0.396	16.2

GRAIN MEAN DM% 84.0

PLOT AREA HARVESTED 0.00260