

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1979

[Full Table of Content](#)



79/R/B/21 Drills and Methods of Applying Fertiliser - Barley

Rothamsted Research

Rothamsted Research (1980) *79/R/B/21 Drills and Methods of Applying Fertiliser - Barley* ; Yields Of The Field Experiments 1979, pp 317 - 318 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/R/B/21

SPRING BARLEY

DRILLS AND METHODS OF APPLYING FERTILISER

Object: To study the effects of different drills and rates and times of applying nitrogen fertiliser on the growth and yield of barley - Bylands.

Sponsor: R. Moffitt.

Design: 3 randomised blocks of 16 plots.

Whole plot dimensions:

DRILLS MF	5.33 x 10.1
DRILLS NIAE	4.27 x 10.1
EXTRA	3.05 x 10.1

Treatments: All combinations of:-

1. DRILLS Drills:
MF 'Massey Ferguson 30' drill, sowing rows 18 cm (7 in) apart
NIAE 'NIAE' drill, sowing rows 18 cm (7 in) apart

2. TOTAL N Total nitrogen fertiliser (kg N):
60
120

3. N METHOD Method of applying nitrogen fertiliser:
CDE Combine drilled at sowing
BCL Broadcast by machine 17 days after sowing
CDE/BCL Half total combine drilled, half broadcast by machine
 17 days after sowing

plus four extra treatments

EXTRA

F60 BCE 'Fiona' drill, sowing rows 15 cm (6 in) apart, 60 kg N
 broadcast by machine at sowing
F120 BCE As previously but using 120 kg N
F60 BCL 'Fiona' drill, 60 kg N broadcast by machine 17 days after sowing
F120 BCL As previously but using 120 kg N

Basal applications: Weedkillers: Bromoxynil and ioxynil (as 'Oxytril CM' at 1.4 kg) with mecoprop at 1.7 kg in 220 l. Fungicide: Tridemorph at 0.53 kg in 220 l.

Seed: Porthos, sown at 160 kg.

Cultivations, etc.: - Subsoiled, tines 100 cm apart and 45 cm deep: 15 Nov, 1978.
Ploughed: 21 Dec. Heavy spring-tine cultivated, rotary harrowed: 27 Apr, 1979.
Seed sown: 30 Apr. Weedkillers applied: 4 June. Fungicide applied: 12 June.
Combine harvested: 29 Aug. Previous crops: Wheat 1977, barley 1978.

NOTES: (1) Observations of growth stages, evenness of growth and wheeling effects were made several times during the season.
(2) Severe grazing by rabbits, and infestations of perennial grasses may have affected yield.

79/R/B/21

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TOTAL N	60	120	MEAN	
DRILLS				
MF	2.85	3.36	3.10	
NIAE	2.89	2.76	2.82	
MEAN	2.87	3.06	2.96	

N METHOD	CDE	BCL	CDE/BCL	MEAN
DRILLS				
MF	3.46	2.91	2.94	3.10
NIAE	2.68	2.70	3.08	2.82
MEAN	3.07	2.81	3.01	2.96

N METHOD	CDE	BCL	CDE/BCL	MEAN
TOTAL N				
60	3.03	2.63	2.95	2.87
120	3.12	2.98	3.07	3.06
MEAN	3.07	2.81	3.01	2.96

DRILLS	N METHOD	CDE	BCL	CDE/BCL
	TOTAL N			
MF	60	3.32	2.72	2.52
	120	3.61	3.11	3.35
NIAE	60	2.74	2.55	3.38
	120	2.63	2.86	2.79

EXTRA	F60 BCE	F120 BCE	F60 BCL	F120 BCL	MEAN
	2.66	3.14	2.48	3.21	2.87

GRAND MEAN 2.94

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	DRILLS	TOTAL N	N METHOD
SED	0.296	0.121	0.121	0.148
TABLE	DRILLS	DRILLS	TOTAL N	DRILLS
	TOTAL N	N METHOD	N METHOD	TOTAL N
				N METHOD
				& EXTRA
SED	0.171	0.209	0.209	0.296

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	30	0.363	12.3

GRAIN MEAN DM% 85.4

PLOT AREA HARVESTED 0.00215