

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1979

[Full Table of Content](#)



79/R/BE/6 N and Pathogen Control - Beans

Rothamsted Research

Rothamsted Research (1980) *79/R/BE/6 N and Pathogen Control - Beans* ; Yields Of The Field Experiments 1979, pp 331 - 332 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/R/BE/6

SPRING BEANS

N AND PATHOGEN CONTROL

Object: To study the effect of enhanced pathogen control on the proportion of nitrogen in the crop derived from the soil, from fertiliser and from nitrogen-fixation. The study was aided by using ^{15}N -labelled fertilisers and spring barley as a crop which did not fix nitrogen - Little Hoos.

Sponsors: J.M. Day, R.J. Roughley, J.F. Witty.

Design: 4 randomised blocks of 12 plots.

Whole plot dimensions: Beans: 3.25 x 4.57, barley 2.13 x 4.57.

Treatments: All combinations of:-

- | | |
|-------------|---|
| 1. PATHCONT | Pathogen control: |
| STANDARD | Standard, pirimicarb foliar spray only |
| ENHANCED | Aldicarb at 10 kg to seedbed plus pirimicarb foliar spray |
| 2. BEANS N | Nitrogen fertiliser (kg N) to beans: |
| 0 | |
| 50 | |
| 100 | |
| 150 | |

plus four extra treatments sown to spring barley and given rates of nitrogen fertiliser (kg N):

BARLEY N

- 0
- 50
- 100
- 150

Standard applications: Barley: Manures: (0:20:20) at 310 kg, combine drilled.
Beans: Insecticide: Pirimicarb at 0.14 kg in 340 l.

Seed: Barley: Porthos, sown at 160 kg.
Beans: Minden, sown at 260 kg.

Cultivations, etc.: - Ploughed: 13 Dec, 1978. Heavy spring-tine cultivated: 21 Apr, 1979. Aldicarb applied, rotary harrowed: 1 May. Barley sown: 3 May. Beans sown: 9 May. Insecticide applied: 12 July. Barley hand harvested: 7 Sept. Beans hand harvested: 24 Sept. Previous cropping: Wheat 1977 and 1978.

NOTES: (1) Content of ^{15}N was assessed in whole plants shortly before harvest.
(2) Nitrogen percentages of grain were measured.

79/R/BE/6

BEANS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

BEANS N PATHCONT	0	50	100	150	MEAN
STANDARD	4.05	4.10	4.25	4.13	4.13
ENHANCED	5.33	5.38	5.86	5.75	5.58
MEAN	4.69	4.74	5.05	4.94	4.86

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PATHCONT	BEANS N	PATHCONT BEANS N
SED	0.131	0.185	0.261

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	21	0.369	7.6

GRAIN MEAN DM% 83.2

PLOT AREA HARVESTED 0.00074

BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

BARLEY N	0	50	100	150	MEAN
	3.48	4.00	4.93	4.72	4.28

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	BARLEY N
SED	0.297

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	9	0.420	9.8

GRAIN MEAN DM% 94.0

PLOT AREA HARVESTED 0.00049