

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1979

[Full Table of Content](#)



79/R/RN/8 Cultivation/WEEDKILLER - Barley

Rothamsted Research

Rothamsted Research (1980) *79/R/RN/8 Cultivation/WEEDKILLER - Barley* ; Yields Of The Field Experiments 1979, pp 76 - 78 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/R/RN/8

CULTIVATION/WEEDKILLER

Object: To study the long-term effects of weedkillers and different methods of primary cultivation on a sequence of crops - Great Harpenden I.

Sponsors: R. Moffitt, G.V. Dyke, J.A. Currie.

The 19th year, barley.

For previous years see 'Details' 1967 and 1973 and 74-78/R/RN/8.

Design: 2 randomised blocks of 12 plots split into 2.

Whole plot dimensions: 12.8 x 15.2.

Treatments: All combinations of:-

Whole plots

1. CULTIVTN Primary cultivations annually:
 PLOUGH Ploughed: 27 Nov, 1978
 ROTAVATE Rotary cultivated by rotary digger: 28 Nov
 DEEPTINE Deep-tine cultivated twice: 27 Nov
2. WEEDCNTL(76) Weed control to beans and potatoes in the rotation beans, wheat, potatoes, barley practised until 1976. Last applied to beans 1976:
 MECHANCL Mechanical
 RESIDUAL Residual weedkiller (duplicated)

Sub plots

3. WEEDKLLR(75) Hormone weedkiller to cereals in the previous rotation, last applied to barley 1975 (basal hormone weedkiller to spring wheat 1977 and barley 1978 and 1979):
 NONE
 HORMONE
4. WEEDKLLR(79) Paraquat weedkiller to cereal stubbles: 23 Oct, 1978
 NONE
 PARAQUAT

NOTE: The combinations of 3 and 4 are tested on half plots: WEEDKLLR(75) NONE, WEEDKLLR(79) NONE and WEEDKLLR(75) HORMONE, WEEDKLLR(79) PARAQUAT on one block, remaining combinations on the other.

- EXTRA plus three extra whole plot treatments:
- SPNGTINE Heavy spring-tine cultivated twice: 27 Nov, 1978. Given simazine to beans 1976, with sub plot tests 3 and 4 above.
 - (SH)PLGH Shallow ploughed: 27 Nov, 1978. Given simazine to beans 1976 and paraquat to cereal stubbles with sub plot test 3 above.

79/R/RN/8

STANDARD Standard cultivations as considered best for each crop.
Ploughed 27 Nov, 1978. Given simazine to beans 1976, with
sub plot tests 3 and 4 above.

NOTE: Paraquat was applied at 0.56 kg ion in 220 l.

Basal applications: Manures: (20:14:14) at 440 kg, combine drilled. Weedkillers:
Bromoxynil and ioxynil (as 'Oxytril CM' at 1.4 kg) and mecoprop at 1.7 kg
in 220 l. Fungicide: Tridemorph at 0.53 kg in 220 l.

Seed: Porthos, sown at 160 kg.

Cultivations, etc.: - Spring-tine cultivated twice, seed sown: 19 Apr, 1979.
Weedkillers applied: 4 June. Fungicide applied: 12 June. Combine harvested:
26 Aug.

EXTRA PLOTS ONLY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

EXTRA	SPNGTINE	(SH)PLGH	STANDARD
WEEDKLLR(75)			
NONE	4.90	4.89	4.78
HORMONE	5.18	5.06	4.74
WEEDKLLR(79)			
NONE	5.01		4.44
PARAQUAT	5.08	4.98	5.08
MEAN	5.04	4.98	4.76

GRAIN MEAN DM% 82.7

SUB PLOT AREA HARVESTED 0.00434

79/R/RN/8

OMITTING EXTRA PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

CULTIVTN	PLOUGH	ROTAVATE	DEEPTINE	MEAN
WEEDCNTL(76)				
MECHANCL	4.80	4.67	4.61	4.70
RESIDUAL	5.04	4.64	4.68	4.79
WEEDKLLR(75)				
NONE	5.03	4.63	4.74	4.80
HORMONE	4.88	4.67	4.58	4.71
WEEDKLLR(79)				
NONE	5.03	4.61	4.62	4.75
PARAQUAT	4.89	4.69	4.70	4.76
MEAN	4.96	4.65	4.66	4.76

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CULTIVTN	WEEDCNTL(76)	WEEDKLLR(75)	WEEDKLLR(79)
SED	0.181	0.156	0.082	0.082

TABLE	CULTIVTN WEEDCNTL(76)	CULTIVTN WEEDKLLR(75)	CULTIVTN WEEDKLLR(79)	
SED	0.313			MIN REP
	0.271	0.206	0.206	MAX-MIN
	0.221			MAX REP

EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:
CULTIVTN 0.141 0.141

WEEDCNTL(76)
MIN REP MECHANCL
MAX-MIN MECHANCL V RESIDUAL
MAX REP RESIDUAL

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	11	0.313	6.6
BLOCK.WP.SP	10	0.245	5.1

GRAIN MEAN DM% 82.8

SUB PLOT AREA HARVESTED 0.00434