

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1980

[Full Table of Content](#)



### 80/W/CS/99 Effects of Breaks on Take -all

#### Rothamsted Research

Rothamsted Research (1981) *80/W/CS/99 Effects of Breaks on Take -all* ; Yields Of The Field Experiments 1980, pp 119 - 121 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/W/CS/99

EFFECTS OF BREAKS ON TAKE-ALL

Object: To study factors affecting the incidence of take-all (*Gaeumannomyces graminis*) and their effects on yields of s. barley - Woburn, Butt Furlong.

Sponsor: D. Hornby.

The eight year, s. barley, s. beans, s. oats.

For previous years see 72/W/CS/99(t) and 73-79/W/CS/99.

Design: 2 randomised blocks of 9 plots, 4 of which are split into 2.

Whole plot dimensions: 5.34 x 15.2.

Treatments: All combinations of:-

Whole plots

1. TREATMNT Crop sequences; soil sterilant and inoculum in 1979:

	1968-71	72	73	74	75	76	77	78	79	80
B 7(S)	B	F	BE	B	B	B	B	B	B(S)	B
B 5	B	B	B	F	BE	B	B	B	B	B
B 4(SI)	B	B	B	B	F	BE	B	B	B(SI)	B
B 3(I)	B	B	B	B	B	F	BE	B	B(I)	B

Sub plots

2. INOCULUM Take-all inoculum:

0	None
I	Inoculated

plus five extra plots testing crop sequences alone:

EXTRA

	1968-71	72	73	74	75	76	77	78	79	80
B 13	B	F	B	B	B	B	B	B	F	BE
B 6	B	B	F	BE	B	B	B	B	B	B
B 1	B	B	B	B	B	B	F	BE	0	B
	B	B	B	B	B	B	B	F	BE	0

B = S. barley, BE = S. beans, 0 = S. oats, F = Fallow  
 (S) = Soil sterilant (1979). Formalin. (I) = Inoculum of take-all (1979) applied on colonised autoclaved oats and combine drilled with barley.  
 I = Inoculum of take-all 1980, applied on colonised autoclaved oats, in the ratio of three oats to one barley seed, broadcast at 310 kg on the surface and rotary cultivated in.

80/W/CS/99

Standard applications:

All crops: Weedkillers: Glyphosate at 1.5 kg in 250 l.

S. barley and s. oats: Manures: (20:10:10) at 450 kg, combine drilled.  
Weedkillers: Mecoprop with bromoxynil and ioxynil as ('Brittox' at 3.5 l) in 280 l with the fungicide. Fungicide: Tridemorph at 0.53 kg.

S. beans: Manures: (0:14:28) at 400 kg. Weedkiller: Simazine at 0.84 kg in 250 l.

Seed: S. barley: Georgie, dressed with ethirimol, sown at 160 kg.

S. oats: Panema, sown at 180 kg.

S. beans: Minden, sown at 200 kg.

Cultivations, etc.:-

All plots: Glyphosate applied: 13 Sept, 1979. Ploughed: 5 Nov.

Spring-tine cultivated with crumbler attached: 28 Feb, 1980.

S. barley: Spring-tine cultivated with crumbler attached: 7 Apr, 1980.

Seed with inoculum treatment applied, all plots rotary cultivated, seed sown: 9 Apr. 'Brittox' with fungicide applied: 13 May. Combine harvested: 21 Aug.

S. oats: Spring-tine cultivated with crumbler attached: 7 Apr, 1980.

Rotary cultivated, seed sown: 9 Apr. 'Brittox' with fungicide applied: 13 May. Combine harvested: 4 Sept.

S. beans: PK applied, seed sown: 28 Feb, 1980. Weedkiller applied: 29 Feb. Combine harvested: 19 Sept.

NOTES: (1) Soil samples were taken before treatments were applied for estimates of *Heterodera avenae* cysts and eggs and incidence of microflora.

(2) Plant samples were taken in July for incidence of take-all.

80/W/CS/99

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT INOCULUM	B 7(S)	B 5	B 4(SI)	B 3(I)	MEAN
0	5.08	5.01	4.75	5.11	4.99
I	4.63	4.96	4.54	4.76	4.72
MEAN	4.85	4.99	4.65	4.94	4.86
EXTRA	B 13 5.11	B 6 5.37	B 1 5.51	MEAN 5.33	

GRAND MEAN 5.06

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	EXTRA	INOCULUM	TREATMNT	INOCULUM TREATMNT
-----				
SED	0.274	0.185	0.274	0.378

EXCEPT WHEN COMPARING MEANS WITH THE SAME LEVEL(S) OF:  
TREATMNT 0.369

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	6	0.274	5.4
BLOCK.WP.SP	10	0.369	7.3

GRAIN MEAN DM% 82.9

SUB PLOT AREA HARVESTED 0.00199