

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1980

[Full Table of Content](#)



80/R/CS/140 Chemical Reference Plots

Rothamsted Research

Rothamsted Research (1981) *80/R/CS/140 Chemical Reference Plots ; Yields Of The Field Experiments 1980*, pp 127 - 130 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of continuous s. barley - Long Hoos V 3.

Sponsors: G.G. Briggs, N. Walker, R. MacDonald.

The sixth year, s. barley.

For previous years see 74-79/R/CS/140.

Design: Single replicate of 32 plots.

Whole plot dimensions: 4.06 x 4.57.

Treatments, applied cumulatively 1974-79 except WEEDKLLR(1) - 1974 and 76 only and WEEDKLLR(2) - 1980 only:
All combinations of:-

1. FUNGCIDE Fungicide:

NONE	None
BENOMYL	Benomyl at 4 kg to the seedbed

2. INSCTCDE Insecticide:

NONE	None
CHLORFEN	Chlorfenvinphos at 2 kg to the seedbed

3. NEMACIDE Nematicide:

NONE	None
ALDICARB	Aldicarb at 6 kg to the seedbed as granules

4. WEEDKLLR(1) Weedkiller in spring:

NONE	None
CHLORTOL	Chlortoluron at 2 kg soon after sowing, applied 1974 and 1976 only

5. WEEDKLLR(2) Weedkiller in autumn:

NONE	None
GLYPHOS	Glyphosate at 1.5 kg to stubble of 1979 s. barley.

NOTE: Glyphosate was applied on 13 Sept, 1979. Other treatments were applied on 5 Mar, 1980.

Basal applications: 'Nitro-Chalk' at 450 kg. Weedkillers: Dicamba with mecoprop and MCPA (as 'Tetralex Plus' at 5.6 l in 340 l).
Fungicide: Tridemorph at 0.53 kg in 340 l.

Seed: Georgie, seed not dressed, sown at 160 kg.

80/R/CS/140

Cultivations, etc.: - Ploughed: 29 Oct, 1979. N applied, spring-tine cultivated: 4 Mar, 1980. Spring-tine cultivated, power harrowed, seed sown: 5 Mar. Weedkillers applied: 11 May. Fungicide applied: 4 June. Combine harvested: 19 Aug.

NOTE: Mildew and aphids were assessed during the season.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

INSCTCDE	NONE	CHLORFEN	MEAN
FUNGCIDE			
NONE	4.68	4.78	4.73
BENOMYL	4.79	4.68	4.73
MEAN	4.74	4.73	4.73
NEMACIDE	NONE	ALDICARB	MEAN
FUNGCIDE			
NONE	4.63	4.84	4.73
BENOMYL	4.77	4.69	4.73
MEAN	4.70	4.76	4.73
NEMACIDE	NONE	ALDICARB	MEAN
INSCTCDE			
NONE	4.74	4.73	4.74
CHLORFEN	4.66	4.80	4.73
MEAN	4.70	4.76	4.73
WEEDKLLR(1)	NONE	CHLORTOL	MEAN
FUNGCIDE			
NONE	4.68	4.78	4.73
BENOMYL	4.62	4.85	4.73
MEAN	4.65	4.82	4.73
WEEDKLLR(1)	NONE	CHLORTOL	MEAN
INSCTCDE			
NONE	4.69	4.78	4.74
CHLORFEN	4.61	4.85	4.73
MEAN	4.65	4.82	4.73
WEEDKLLR(1)	NONE	CHLORTOL	MEAN
NEMACIDE			
NONE	4.53	4.87	4.70
ALDICARB	4.77	4.76	4.76
MEAN	4.65	4.82	4.73

80/R/CS/140

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

WEEDKLLR(2)	NONE	GLYPHOS	MEAN
FUNGCIDE			
NONE	4.66	4.80	4.73
BENOMYL	4.65	4.82	4.73
MEAN	4.65	4.81	4.73
WEEDKLLR(2)	NONE	GLYPHOS	MEAN
INSCTCDE			
NONE	4.67	4.80	4.74
CHLORFEN	4.64	4.82	4.73
MEAN	4.65	4.81	4.73
WEEDKLLR(2)	NONE	GLYPHOS	MEAN
NEMACIDE			
NONE	4.64	4.77	4.70
ALDICARB	4.67	4.86	4.76
MEAN	4.65	4.81	4.73
WEEDKLLR(2)	NONE	GLYPHOS	MEAN
WEEDKLLR(1)			
NONE	4.60	4.70	4.65
CHLORTOL	4.71	4.92	4.82
MEAN	4.65	4.81	4.73
INSCTCDE	NONE	CHLORFEN	
NEMACIDE	NONE	ALDICARB	NONE ALDICARB
FUNGCIDE			
NONE	4.64	4.73	4.62
BENOMYL	4.85	4.73	4.70
			4.95
INSCTCDE	NONE	CHLORFEN	
WEEDKLLR(1)	NONE	CHLORTOL	NONE CHLORTOL
FUNGCIDE			
NONE	4.67	4.70	4.70
BENOMYL	4.71	4.87	4.52
			4.87
NEMACIDE	NONE	ALDICARB	
WEEDKLLR(1)	NONE	CHLORTOL	NONE CHLORTOL
FUNGCIDE			
NONE	4.50	4.75	4.86
BENOMYL	4.56	4.99	4.67
			4.82
NEMACIDE	NONE	ALDICARB	
WEEDKLLR(1)	NONE	CHLORTOL	NONE CHLORTOL
INSCTCDE			
NONE	4.66	4.83	4.72
CHLORFEN	4.41	4.91	4.81
			4.74
			4.79

80/R/CS/140

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

INSCTCDE	NONE	CHLORFEN		
WEEDKLLR(2)	NONE	GLYPHOS	NONE	GLYPHOS
FUNGCIDE				
NONE	4.64	4.73	4.69	4.88
BENOMYL	4.70	4.87	4.59	4.76
NEMACIDE	NONE	ALDICARB		
WEEDKLLR(2)	NONE	GLYPHOS	NONE	GLYPHOS
FUNGCIDE				
NONE	4.55	4.71	4.78	4.90
BENOMYL	4.73	4.82	4.57	4.81
NEMACIDE	NONE	ALDICARB		
WEEDKLLR(2)	NONE	GLYPHOS	NONE	GLYPHOS
INSCTCDE				
NONE	4.72	4.77	4.62	4.83
CHLORFEN	4.55	4.76	4.72	4.88
WEEDKLLR(1)	NONE	CHLORTOL		
WEEDKLLR(2)	NONE	GLYPHOS	NONE	GLYPHOS
FUNGCIDE				
NONE	4.58	4.78	4.74	4.82
BENOMYL	4.61	4.62	4.68	5.01
WEEDKLLR(1)	NONE	CHLORTOL		
WEEDKLLR(2)	NONE	GLYPHOS	NONE	GLYPHOS
INSCTCDE				
NONE	4.68	4.70	4.66	4.90
CHLORFEN	4.51	4.71	4.76	4.93
WEEDKLLR(1)	NONE	CHLORTOL		
WEEDKLLR(2)	NONE	GLYPHOS	NONE	GLYPHOS
NEMACIDE				
NONE	4.44	4.62	4.83	4.91
ALDICARB	4.75	4.79	4.60	4.92

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

ONE FACTOR TABLES 0.101
TWO FACTOR TABLES 0.143
THREE FACTOR TABLES 0.202

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	6	0.286	6.0

GRAIN MEAN DM% 82.5

PLOT AREA HARVESTED 0.00075