

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/R/CS/248 Direct Drilling and Slug Control

Rothamsted Research

Rothamsted Research (1981) *80/R/CS/248 Direct Drilling and Slug Control* ; Yields Of The Field Experiments 1980, pp 190 - 191 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/CS/248

DIRECT DRILLING AND SLUG CONTROL

Object: To study the effects of a range of materials and methods of application on the control of slugs and on the yield of direct-drilled w. wheat - Pastures.

Sponsors: G.C. Scott, D.C. Griffiths, C.A. Edwards.

The first year, w. wheat.

Design: 4 randomised blocks of 11 plots.

Whole plot dimensions: 5.33 x 7.62.

Treatments:

TREATMNT	Materials, methods and rates of application:
NONE	None (duplicated)
MB SD	Methiocarb seed dressing (0.2% weight of seed)
SAN 1 SD	'SAN 155' seed dressing (0.1% weight of seed)
SAN 3 SD	'SAN 329' seed dressing (0.1% weight of seed)
MB 1 BCE	Methiocarb at 5.6 kg broadcast 7 Nov before drilling
MB 1 BCL	Methiocarb at 5.6 kg broadcast 19 Nov after drilling
MD CD	Metaldhyde at 7.8 kg combine drilled
MB 1 CD	Methiocarb at 5.6 kg combine drilled
MB 2 CD	Methiocarb at 11.2 kg combine drilled
IOX S	Ioxynil spray (as 'Totril' at 2.8 l) in 450 l before drilling

Basal applications: Manures: 'Nitro-Chalk' at 190 kg and later at 450 kg. Weedkillers: Diquat at 0.59 kg ion in 220 l. Mecoprop at 2.5 l in 220 l. Glyphosate at 2.1 l in 220 l. Isoproturon at 2.9 kg in 250 l. Insecticide: Demeton-s-methyl at 0.24 kg in 250 l.

Seed: Flanders, sown at 180 kg.

Cultivations, etc.: - Diquat applied: 2 Oct, 1979. Mecoprop applied: 9 Oct. Glyphosate applied: 20 Oct. Seed sown: 13 Nov. 'Nitro-Chalk' applied at 190 kg: 18 Feb, 1980. 'Nitro-Chalk' applied at 450 kg: 15 Apr. Isoproturon applied: 24 Apr. Insecticide applied: 24 June. Combine harvested: 21 Aug. Previous crops: Clover 1978 and 1979.

NOTE: Slug counts were made before drilling and during the autumn. Plants were counted and scored for slug damage in Dec, Jan and Apr.

80/R/CS/248

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
NONE	6.82
MB SD	6.93
SAN 1 SD	6.64
SAN 3 SD	6.68
MB 1 BCE	7.28
MB 1 BCL	6.97
MD CD	6.16
MB 1 CD	5.85
MB 2 CD	5.17
IOX S	6.57
MEAN	6.54

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT
-----	-----
SED	0.396 MIN REP
	0.343 MAX-MIN

TREATMNT
 MAX-MIN NONE V ANY OF REMAINDER
 MIN REP ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	31	0.560	8.6
GRAIN MEAN DM%	83.4		
PLOT AREA HARVESTED	0.00213		