

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/R/CS/250 Control of Cephalosporium

Rothamsted Research

Rothamsted Research (1981) *80/R/CS/250 Control of Cephalosporium ; Yields Of The Field Experiments 1980*, pp 192 - 193 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/CS/250

CONTROL OF CEPHALOSPORIUM

Object: To study the effects of aldicarb, benomyl and HCH on the incidence of Cephalosporium Stripe (*Cephalosporium gramineum*) and on the yield of w. wheat - New Zealand.

Sponsors: R.J. Gutteridge, K.E. Fletcher.

The first year, w. wheat.

Design: 3 randomised blocks of 8 plots.

Whole plot dimensions: 2.67 x 6.10.

Treatments:

CHEMICAL	Chemicals and methods of application:
NONE	None (5 plots per block)
ALDICARB	Aldicarb at 10 kg worked in to seedbed
BENOMYL	Benomyl at 20 kg worked in to seedbed
HCH	HCH seed dressing at 0.4 g per kg seed

NOTE: (1) Soil treatments were applied on 8 Oct, 1979.
(2) Benomyl was applied as a drench in 11,200 l of water.

Basal applications: Manures: (10:23:23) at 250 kg, combine drilled.
'Nitro-Chalk' at 340 kg. Weedkiller: Mecoprop at 2.5 l in 220 l.
Growth regulator: Chlormequat at 1.4 l in 250 l.

Seed: Bounty, sown at 180 kg.

Cultivations, etc.: - Chisel ploughed: 29 June, 1979, 3 July, 4 July, 17 July. Rotary cultivated: 6 July, 20 Aug. Spring-tine cultivated: 17 July, 27 Sept. Rotary harrowed, seed sown: 8 Oct. Weedkiller applied: 1 Mar, 1980. N applied: 8 Apr. Growth regulator applied: 6 May. Combine harvested: 23 Aug. Previous crops: Grass 1978 and 1979.

NOTES: Wireworm populations were assessed in August 1979, plant emergence in November, plant and shoot damage by stem borers in February, wireworms in May and incidence of Cephalosporium, take-all, and Phialophora in June.

80/R/CS/250

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

CHEMICAL	NONE	ALDICARB	BENOMYL	HCH	MEAN
	9.30	9.30	9.41	9.14	9.29

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHEMICAL
-----	-----
SED	0.172 MIN REP 0.134 MAX-MIN

	CHEMICAL
MAX-MIN	NONE V ANY OF THE REMAINDER
MIN REP	ANY OF THE REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	18	0.211	2.3

GRAIN MEAN DM% 85.3

PLOT AREA HARVESTED 0.00161