Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1980



Full Table of Content

80/S/CS/1 Fungicides, N and Growth Regulator

Rothamsted Research

Rothamsted Research (1981) 80/S/CS/1 Fungicides, N and Growth Regulator; Yields Of The Field Experiments 1980, pp 197 - 201 - DOI: https://doi.org/10.23637/ERADOC-1-31

FUNGICIDES, N AND GROWTH REGULATOR

Object: To study the effects of fungicides and of a growth regulator, and rates and times of applying nitrogen fertiliser on the incidence of foliar diseases and on the yield of w. barley - Saxmundham, Oldershaw's and Garner's plots.

Sponsors: F.V. Widdowson, J.F. Jenkyn, A. Penny.

The 15th year, w. barley.

For previous years see 66/C/30(t), 67/C/23(t), 68/C/39, 69-70/S/CS/1, 71/S/CS/1(t), 72/S/CS/1(t) and 73-79/S/CS/1.

Design: A single replicate of 24 plots split into 2, arranged as 2 blocks of 12 plots split into 2. Treatments to w. wheat 1966-1976 and to w. and s. barley 1977-1979 have been ignored.

Whole plot dimensions: 2.44 x 40.2.

Treatments: All combinations of:-

Whole plots

1. MILDFUNG Fungicide to control mildew: Tridemorph at 0.53 kg in 280 l.

NONE None

SPRAYED Sprayed 23 Apr, 1980 and 14 May

2. E N RATE Rates of early spring nitrogen fertiliser (kg N):

105 140

3. E N TIME Times of applying early spring nitrogen fertiliser:

FEB+MAR 35 kg on 12 Feb remainder on 25 Mar MAR All on 25 Mar

4. APR N GR Nitrogen fertiliser in April (kg N) and growth regulator:

0 0 None 35 0 35 on 23 Apr

35 GR 35 on 23 Apr plus mepiquat chloride + ethephon ('Terpal' at 2.46 1) in 280 1 on 23 Apr

Half plots

5. EYESFUNG Fungicide to control eyespot: Carbendazim (as 'Bavistin' at 0.50 kg) in 280 l

NONE None SPRAYED Sprayed 23 Apr

Basal applications: Manures: N, P205 & K20 each at 51 kg as (15:15:15). Weedkillers: Chlortoluron at 5.6 kg in 220 l in autumn. 'Wheatclene' (1.26 kg of solid (metoxuron and simazine) plus 1.26 l of liquid (barban)) mixed with ioxynil at 1.3 kg plus mecoprop at 1.9 kg in 220 1.

Seed: Sonja, sown at 180 kg.

Cultivations, etc.:- Seed sown, NPK applied: 26 Sept, 1979. Chlortoluron applied: 27 Sept. Spring weedkillers applied: 31 Mar, 1980. Combine harvested: 29 July.

NOTE: Straw heights were measured in late June. N content of grain was GRAIN TONNES/HECTARE

**** TABLES OF MEANS ****

E N RATE	105	140	MEAN	
NONE	8.31	8.45	8.38	
SPRAYED	8.23	8.75	8.49	
MEAN	8.27	8.60	8.44	
E N TIME MILDFUNG	FEB+MAR	MAR	MEAN	
NONE	8.27	8.49	8.38	
SPRAYED	8.41	8.58	8.49	
MEAN	8.34	8.53	8.44	
E N TIME E N RATE	FEB+MAR	MAR	MEAN	
105	8.16	8.38	8.27	
140	8.52	8.68	8.60	
MEAN	8.34	8.53	8.44	
APR N GR MILDFUNG	0 0	35 0	35 GR	MEAN
NONE	8.13	8.40	8.61	8.38
SPRAYED	7.99	8.61	8.87	8.49
MEAN	8.06	8.51	8.74	8.44
APR N GR E N RATE	0 0	35 0	35 GR	MEAN
105	7.79	8.47	8.54	8.27
140	8.33	8.54	8.93	8.60
MEAN	8.06	8.51	8.74	8.44

GRAIN TONNES/HECTARE

**** TABLES OF MEANS ****

***	* TABLES OF	MEANS ***	**				
	APR N GR	0 0	35 0	35 GR	MEAN		
	E N TIME						
	FEB+MAR	8.00	8.37	8.64	8.34		
	MAR	8.13	8.64	8.84	8.53		
	MEAN	8.06	8.51	8.74	8.44		
	EYESFUNG	NONE	SPRAYED	MEAN			
	MILDFUNG						
	NONE	8.44	8.32	8.38			
	SPRAYED	8.38	8.60	8.49			
	MEAN	8.41	8.46	8.44			
	EYESFUNG	NONE	SPRAYED	MEAN			
	E N RATE						
	105	8.22	8.32	8.27			
	140	8.60	8.60	8.60			
	MEAN	8.41	8.46	8.44			
	EYESFUNG	NONE	SPRAYED	MEAN			
	E N TIME	0.21	0.26	0.24			
	FEB+MAR	8.31	8.36	8.34			
	MAR	8.50	8.56	8.53			
	MEAN	8.41	8.46	8.44			
	EYESFUNG APR N GR	NONE	SPRAYED	MEAN			
	0 0	8.02	8.11	8.06			
	35 0	8.53	8.48				
	35 GR		8.80				
	35 GK	8.68	0.00	0.74			
	MEAN	8.41	8.46	8.44			
	E N RATE	105		140			
	E N TIME MILDFUNG	FEB+MAR	MAR	The second secon	MAR		
	NONE	8.14	8.48	8.40	8.50		
	SPRAYED	8.18	8.28	8.63	8.87		
	E N RATE	105			140		
	APR N GR	0 0	35 0	35 GR	0 0	35 0	35 GR
	MILDFUNG						
	NONE	7.95	8.48	8.50	8.32	8.32	8.71
	SPRAYED	7.63	8.47	8.59	8.35	8.76	9.15
	E N TIME	FEB+MAR			MAR		
	APR N GR	0 0	35 0	35 GR	0 0	35 0	35 GR
	MILDFUNG	8.11	8.18	8.51	8.16	8.62	8.70
	SPRAYED	7.88	8.56	8.77	8.10	8.67	8.97

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

INDEES OF	FILANO					
E N TIME	FEB+MAR			MAR		
APR N GR E N RATE	0 0	35 0	35 GR	0 0	35 0	35 GR
105	7.81	8.31	8.35	7.78	8.64	8.74
140	8.19	8.43	8.93	8.48	8.64	8.94
E N RATE	105		140			
EYESFUNG MILDFUNG	NONE	SPRAYED	NONE	SPRAYED		
NONE	8.25	8.37	8.62	8.27		
SPRAYED	8.20	8.26	8.57	8.93		
E N TIME	FEB+MAR	20.00	MAR			
EYESFUNG MILDFUNG	NONE	SPRAYED	NONE	SPRAYED		
NONE	8.37	8.17	8.50	8.48		
SPRAYED	8.26	8.55	8.51	8.65		
E N TIME	FEB+MAR	182.5	MAR			
EYESFUNG E N RATE	NONE	SPRAYED	NONE	SPRAYED		
105	8.08	8.23	8.36	8.41		
140	8.54	8.49	8.65	8.72		
APR N GR	0 0		35 0		35 GR	
EYESFUNG MILDFUNG	NONE	SPRAYED	NONE	SPRAYED	NONE	SPRAYED
NONE	8.16	8.11	8.47	8.33	8.68	8.53
SPRAYED	7.88	8.10	8.59	8.63	8.68	9.06
APR N GR	0 0	16.8	35 0	8 35.	35 GR	
EYESFUNG E N RATE	NONE	SPRAYED	NONE	SPRAYED	NONE	SPRAYED
105	7.66	7.92	8.40	8.55	8.60	8.49
140	8.38	8.29	8.65	8.42	8.76	9.10
APR N GR	0 0		35 0		35 GR	
EYESFUNG E N TIME	NONE	SPRAYED	NONE	SPRAYED	NONE	SPRAYED
FEB+MAR	7.97	8.03	8.49	8.25	8.48	8.80
MAR	8.07	8.18	8.56	8.72	8.88	

GRAIN TONNES/HECTARE

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	EYESFUNG	MILDFUNG* EYESFUNG	E N RATE* EYESFUNG	E N TIME* EYESFUNG
SED	0.091	0.129	0.129	0.129
TABLE	APR N GR* EYESFUNG	MILDFUNG* E N RATE EYESFUNG	MILDFUNG* E N TIME EYESFUNG	E N RATE* E N TIME EYESFUNG
SED	0.158	0.182	0.182	0.182
TABLE	MILDFUNG* APR N GR EYESFUNG	E N RATE* APR N GR EYESFUNG	E N TIME* APR N GR EYESFUNG	
SED	0.223	0.223	0.223	

^{*} USE ONLY TO COMPARE THE TWO LEVELS OF EYESFUNG FOR THE SAME LEVEL(S) OF THE OTHER FACTOR(S)

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

 STRATUM
 DF
 SE
 CV%

 BLOCK.WP.SP
 9
 0.316
 3.7

GRAIN MEAN DM% 84.7

SUB PLOT AREA HARVESTED 0.00380