

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)

Her Majesty's Stationery Office
LONDON
1981

This document is available to members of the Scientific Community and other qualified persons only. It is not to be distributed outside the United Kingdom.

Printed in Great Britain

80/R/WW/7 Winter Wheat Fungicides and Soil-borne Diseases

Rothamsted Research

Rothamsted Research (1981) *80/R/WW/7 Winter Wheat Fungicides and Soil-borne Diseases ; Yields Of The Field Experiments 1980*, pp 238 - 239 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/WW/7

WINTER WHEAT

FUNGICIDES AND SOIL-BORNE DISEASES

Object: To study the effects of a range of soil-applied fungicides on soil-borne diseases and yield of w. wheat - Claycroft.

Sponsor: G.L. Bateman.

Design: 4 randomised blocks of 7 plots.

Whole plot dimensions: 2.13 x 12.2.

Treatments:

FUNGICIDE	Fungicides and times and methods of application:
NONE	None (duplicated)
BEN A	Benomyl at 20 kg, worked in to seedbed in autumn
BEN S	Benomyl at 20 kg, as a drench in 5400 l in spring
CHL A	Chloroneb at 14 kg, worked in to seedbed in autumn
KWG A+S	'KWG 0599' at 23 kg a.i. worked in to seedbed in autumn, repeated as a drench in 5400 l in spring
NUA A+S	Nuarimol at 4.4 kg, worked in to seedbed in autumn, repeated as a drench in 5400 l in spring

Basal applications: Manures: (10:23:23) at 250 kg, combine drilled.
'Nitro-Chalk' at 560 kg. Weedkillers: Paraquat at 0.56 kg ion in 220 l.
Mecoprop at 2.5 l in 280 l.

Seed: Flanders, no seed dressing, sown at 190 kg.

Cultivations, etc.: - Heavy spring-tine cultivated twice: 20 Sept, 1979.
Heavy spring-tine cultivated: 4 Oct. Autumn treatments applied: 11 Oct.
Paraquat applied: 16 Oct. Seed sown: 18 Oct. N applied: 11 Apr, 1980.
Mecoprop applied: 16 Apr. Spring treatments applied: 24 Apr.
Combine harvested: 21 Aug. Previous crops: W. wheat 1978, s. barley 1979.

NOTE: Foot and root diseases were assessed monthly between April and July.

80/R/WW/7

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE	NONE	BEN A	BEN S	CHL A	KWG A+S	NUA A+S	MEAN
	9.47	9.60	9.78	9.42	9.75	8.65	9.45

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGCIDE
SED	0.207 MIN REP 0.179 MAX-MIN

FUNGCIDE
MAX-MIN NONE V ANY OF REMAINDER
MIN REP ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	19	0.292	3.1

GRAIN MEAN DM% 83.4

PLOT AREA HARVESTED 0.00176