

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/R/B/8 and 80/W/B/8 Spring Barley Physiological Study on Contrasted Sites

Rothamsted Research

Rothamsted Research (1981) *80/R/B/8 and 80/W/B/8 Spring Barley Physiological Study on Contrasted Sites* ; Yields Of The Field Experiments 1980, pp 269 - 271 - DOI:

<https://doi.org/10.23637/ERADOC-1-31>

80/R/B/8 and 80/W/B/8

SPRING BARLEY

PHYSIOLOGICAL STUDY ON CONTRASTED SITES

Object: To study the effects of a range of nitrogen fertiliser dressings and two contrasted sites on the grain physiology and yield of two varieties of s. barley - Rothamsted (R) Delafield and Woburn (W) Lansome III.

Sponsor: J.N. Gallagher.

Design: 3 randomised blocks of 12 plots.

Whole plot dimensions: 3.0 x 16.2.

Treatments: All combinations of:-

1. VARIETY Varieties:

ARKROYAL	Ark Royal
PORTHOS	Porthos

2. N Rates of nitrogen fertiliser (kg N):

R	W
0	0
25	30
50	60
75	90
100	120
125	150

NOTE: Ark Royal was sown at 140 kg and Porthos at 160 kg. These seed rates each gave 350 seeds per square metre.

Basal applications:

Delafield (R): Manures: (0:20:20) at 310 kg. Weedkillers: Dicamba with mecoprop and MCPA ('Banlene Plus' at 5.0 l) in 250 l applied with the fungicide. Fungicide: Tridemorph at 0.53 kg.

Lansome III (W): Manures: FYM at 50 t, (0:20:20) at 310 kg. Weedkillers Mecoprop with bromoxynil and ioxynil ('Brittox' at 3.5 l) in 280 l applied with the tridemorph. Fungicides: Tridemorph at 0.53 kg; triadimefon 0.13 kg in 220 l applied with the insecticide. Insecticide: Demeton-s-methyl at 0.24 l.

Cultivations, etc.:-

Delafield (R): Subsoiled, tines 160 cm apart and 40 cm deep: 12 Nov, 1979. Chisel ploughed twice: 3 Dec, 4 Dec. Spring-tine cultivated: 6 Apr, 1980. PK and N applied: 7 Apr. Spring-tine cultivated, seed sown: 8 Apr. Weedkiller and fungicide applied: 27 May. Combine harvested: 1 Sept. Previous crops: Beans and kale 1978, potatoes 1979.

80/R/B/8 and 80/W/B/8

Lansome III (W): Subsoiled, tines 140 cm apart and 50 cm deep: 12 Aug, 1979. FYM applied, ploughed: 21 Aug. Heavy spring-tine cultivated: 29 Feb, 1980. PK applied: 13 Mar. Heavy spring-tine cultivated: 4 Apr. N applied, spring-tine cultivated with crumbler attached: 5 Apr. Seed sown: 10 Apr. Weedkiller with tridemorph applied: 30 May. Triadimefon with insecticide applied: 3 July. Combine harvested: 28 Aug. Previous crops: S. barley 1978, grass 1979.

- NOTES: (1) Plant populations were counted at Rothamsted and Woburn and crop cover and growth were assessed at Woburn.
 (2) Because of rabbit damage the yield from one plot at Rothamsted was lost with treatment combination, VARIETY ARKROYAL N 50. An estimated value was used in the analysis.

80/R/B/8 DELAFIELD (R)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

N	0	25	50	75	100	125	MEAN
VARIETY							
ARKROYAL	4.53	5.60	5.65	6.65	6.63	6.27	5.89
PORTHOS	4.83	5.39	6.28	6.37	6.63	6.97	6.08
MEAN	4.68	5.50	5.96	6.51	6.63	6.62	5.98

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	N	VARIETY N

SED	0.105	0.183	0.258

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	21	0.316	5.3

GRAIN MEAN DM% 77.1

PLOT AREA HARVESTED 0.00229

80/W/B/8 LANSOME III (W)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

N	0	30	60	90	120	150	MEAN
VARIETY							
ARKROYAL	4.07	5.01	6.42	6.49	7.23	7.80	6.17
PORTHOS	4.02	4.61	5.99	6.30	6.88	6.90	5.78
MEAN	4.04	4.81	6.20	6.40	7.06	7.35	5.98

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	N	VARIETY N

SED	0.177	0.307	0.434

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	22	0.532	8.9
GRAIN MEAN DM%	77.4		
PLOT AREA HARVESTED	0.00228		