

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)

Her Majesty's Stationery Office
LONDON
Printed in Great Britain
1981

80/R/BE/5 Spring Beans Effects of Pest and Pathogen Control

Rothamsted Research

Rothamsted Research (1981) *80/R/BE/5 Spring Beans Effects of Pest and Pathogen Control* ; Yields Of The Field Experiments 1980, pp 283 - 284 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/BE/5

SPRING BEANS

EFFECTS OF PEST AND PATHOGEN CONTROL

Object: To assess the benefits from three amounts of pest and disease control on irrigated and unirrigated s. beans - Gt Knott II.

Sponsors: J. McEwen, R. Bardner, A.J. Cockbain, J.M. Day, K.E. Fletcher, B.J. Legg, G.A. Salt, R.M. Webb, J.F. Witty, D.P. Yeoman.

Design: 4 randomised blocks of 2 plots split into 3.

Whole plot dimensions: 4.27 x 13.7.

Treatments: All combinations of:-

Whole plots

1. IRRIGATN	Irrigation:
NONE	None
FULL	Full (total 124 mm)

Sub plots

2. PATHCONT	Pest and pathogen control
STANDARD	None
ENHANCED	Permethrin at 0.10 kg on 23 Apr, 1980 Pirimicarb at 0.14 kg on 2 June Benomyl at 0.56 kg on 16 July
FULL	Aldicarb at 10 kg on 3 Mar Permethrin at 0.10 kg on 23 Apr Aluminium tris-ethyl phosphonate at 2.0 kg on 23 Apr Pirimicarb at 0.14 kg on 2 June Benomyl at 0.56 kg on 16 July Benomyl at 0.56 kg on 18 Aug

- NOTES: (1) A planned application of pirimicarb to all PATH CONT treatments to control black fly (*Aphis fabae*) was not applied as numbers of this pest were few.
- (2) Irrigation was applied to reduce a deficit of 50 mm to 25 mm before pod set, and from 80 mm to 55 mm after pod set. (mm water):

17 May	9
19 May	21
24 May	25
27 May	20
5 June	6
6 June	6
8 June	12
13 June	25

124

80/R/BE/5

(3) Treatment sprays were applied in 340 l.

Basal applications: Manures: Chalk at 3 t. Weedkillers: Trietazine with simazine (as 'Remtal SC' at 2.5 l) in 250 l.

Seed: Minden, sown at 150 kg. (500,000 seeds per hectare).

Cultivations, etc.: - Chalk applied: 8 Oct, 1979. Ploughed: 29 Nov. Spring-tine cultivated: 1 Mar, 1980. Spring-tine cultivated, rotary harrowed: 3 Mar. Seed sown: 4 Mar. Weedkillers applied: 21 Mar. Combine harvested: 18 Sept. Previous crops: W. wheat 1978, s. barley 1979.

NOTE: Plant counts were made after establishment and components of yield measured before harvest. Total above-ground dry matter and N content were measured in August. Migratory nematodes, root and foliar fungi, aphids, weevils and viruses were counted at intervals during the season. N content of grain was measured.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PATHCONT IRRIGATN	STANDARD	ENHANCED	FULL	MEAN
NONE	3.91	4.64	5.63	4.72
FULL	3.64	4.26	4.88	4.26
MEAN	3.77	4.45	5.25	4.49

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PATHCONT	IRRIGATN* PATHCONT

SED	0.186	0.263

* WITHIN THE SAME LEVEL OF IRRIGATN ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	12	0.372	8.3

GRAIN MEAN DM% 78.4

SUB PLOT AREA HARVESTED 0.00293