

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1980

[Full Table of Content](#)



## 80/R/BE/9 Spring Beans Fungicides

### Rothamsted Research

Rothamsted Research (1981) *80/R/BE/9 Spring Beans Fungicides ; Yields Of The Field Experiments 1980*, pp 288 - 289 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/R/BE/9

SPRING BEANS

FUNGICIDES

Object: To study the effects of fungicides and methods of application on the incidence of diseases and on the yield of s. beans - Long Hoos III 7.

Sponsors: G.A. Salt, J. McEwen, D.P. Yeoman.

Design: 2 randomised blocks of 20 plots.

Whole plot dimensions: 2.03 x 2.13.

Treatments: All combinations of:-

1. AL METH            Methods of applying aluminium tris (ethyl phosphonate) ('Aliette')(all rates are of a.i.):

NONE	Not applied
SEEDRESS	Seed dressing at 4.0 g per kg seed
SPRAY E	Early foliar spray at 3.4 kg
SPRAY L	Late foliar spray at 3.4 kg

2. BEN METH            Methods of applying benomyl:

NONE	Not applied
SEEDRESS	Seed dressing at 6.0 g per kg seed
SPRAY E	Early foliar spray at 1.0 kg
SPRAY L	Late foliar spray at 1.0 kg

plus four extra treatments:

EXTRA

NONE	No fungicides
STICKER	Methyl cellulose sticker as used for all seed dressing treatments
BE TH SD	Benomyl + thiram seed dressing (duplicated) at 2.0 g of each material per kg seed

NOTE: (1) Seed was sown by hand in rows 51 cm apart, seed spaced 5 cm apart in the row.

(2) Early foliar sprays were applied on 28 May in 700 l and late sprays on 22 July in 1200 l water.

Basal applications: Weedkillers: Trietazine and simazine (as 'Remtal SC' at 3.0 l) in 340 l. Insecticides: Permethrin at 0.15 kg in 340 l. Pirimicarb at 0.14 kg in 340 l.

Seed: Minden.

Cultivations, etc.:- Ploughed: 14 Nov, 1979. Spring-tine cultivated: 7 Apr, 1980. Power harrowed, seed sown: 8 Apr. Weedkillers applied: 15 Apr. Permethrin applied: 23 May. Pirimicarb applied: 11 July. Hand harvested: 7 Oct. Previous crops: Potatoes 1978, w. wheat 1979.

80/R/BE/9

NOTE: Emergence counts were made in early June, and root diseases were assessed in late July.

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

BEN METH AL METH NONE	NONE	SEEDRESS	SPRAY E	SPRAY L	MEAN
NONE	4.17	4.65	4.07	5.41	4.58
SEEDRESS	4.67	5.06	4.91	4.98	4.90
SPRAY E	4.75	4.97	4.43	5.04	4.80
SPRAY L	3.61	5.14	5.15	4.85	4.68
MEAN	4.30	4.95	4.64	5.07	4.74

  

EXTRA	NONE	STICKER	BE TH SD	MEAN
	4.15	5.02	5.11	4.85

GRAND MEAN 4.76

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	EXTRA	AL METH	BEN METH	AL METH BEN METH & EXTRA	MIN REP MAX-MIN
SED	0.509 0.402	0.255	0.255	0.509 0.402	

EXTRA  
MAX-MIN BE TH SD V ANY OF THE REMAINDER  
MIN REP ANY OF THE REMAINDER

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	20	0.509	10.7
GRAIN MEAN DM%	87.3		
PLOT AREA HARVESTED	0.00015		