Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1980



Full Table of Content

80/R/P/7 Potatoes Varieties and Times of Applying Fungicides

Rothamsted Research

Rothamsted Research (1981) 80/R/P/7 Potatoes Varieties and Times of Applying Fungicides; Yields Of The Field Experiments 1980, pp 333 - 334 - DOI: https://doi.org/10.23637/ERADOC-1-31

80/R/P/7

POTATOES

VARIETIES AND TIMES OF APPLYING FUNGICIDES

Object: To study the effects of two times of applying two fungicides to seed tubers of three varieties on the yield and infection of progeny tubers and on disease incidence on the stored produce - Sawyers II.

Sponsors: G.A. Hide, G.R. Cayley.

Design: 4 randomised blocks of 15 plots.

Whole plot dimensions: 1.42 x 9.53.

Treatments: All combinations of:-

FUNGCIDE Fungicides and times of application:

NONE None

IMAZ L Imazalil at lifting in 1979

IMAZ L+B Imazalil at lifting in 1979 and at boxing

THIA L Thiabendazole at lifting in 1979

THIA L+B Thiabendazole at lifting in 1979 and at boxing

VARIETY Varieties:

CROWN Pentland Crown
DESIREE Desiree
EDWARD King Edward

NOTES: (1) Lifting treatments were applied on the following dates: Pentland Crown 22 Oct, 1979, Desiree 12 Sept, King Edward 13 Sept. Boxing treatments were all done on 13 Feb 1980.

(2) Fungicides were applied at 18 g per tonne of seed by dipping tubers in a 0.2% solution for five minutes.

Basal applications: Manures: Chalk at 7.5 t. (0:20:20) at 1260 kg. FYM at 25 t. Epsom salts at 145 kg mixed with kieserite at 145 kg. (13:13:20) at 1510 kg. Weedkillers: Linuron at 1.1 kg in 250 l. Fungicide: Mancozeb at 1.3 kg in 250 l applied six times, with insecticide on the third and fourth occasions. Insecticide: Pirimicarb at 0.14 kg. Desiccant: BOV at 170 l.

Cultivations, etc.:- Heavy spring-tine cultivated: 17 Sept, 1979. Chalk applied: 9 Oct. Subsoiled with tines 160 cm apart and 40 cm deep: 1 Nov. PK applied: 13 Nov. FYM applied 21 Dec. Epsom salts and kieserite applied: 31 Dec. Ploughed: 8 Jan, 1980. Heavy spring-tine cultivated: 14 Apr. NPK applied, spike rotary cultivated, ridged: 15 Apr. Hand planted, ridges split back: 16 Apr. Grubbed: 24 Apr. Rotary ridged: 28 Apr. Weedkillers applied: 18 May. Fungicide applied: 18 June, 30 June, 11 July, 24 July, 5 Aug, 18 Aug. Insecticide applied: 11 July, 24 July. Desiccant applied: 19 Sept. Lifted: 3 Oct. Previous crops: W. oats 1978, w. wheat 1979.

NOTE: Emergence counts were made in late May, and plant and stem counts before harvest. Disease assessments were made on harvested produce.

80/R/P/7

TOTAL TUBERS TONNES/HECTARE

**** TABLES OF MEANS ****

VARIETY	CROWN	DESIREE	EDWARD	MEAN
FUNGCIDE		HAT		
NONE	70.8	62.8	59.3	64.3
IMAZ L	69.4	59.4	60.2	63.0
IMAZ L+B	60.0	61.9	54.4	58.8
THIA L	69.3	59.6	59.7	62.9
THIA L+B	72.2	63.0	58.6	64.6
MEAN	68.3	61.3	58.4	62.7

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	FUNGCIDE	VARIETY	FUNGCIDE VARIETY	
SED	1.62	1.25	2.80	

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	42	3.96	6.3

PERCENTAGE WARE 4.44CM (1.75 INCH) RIDDLE

***** TABLES OF MEANS *****

VARIETY	CROWN	DESIREE	EDWARD	MEAN
FUNGCIDE				
NONE	89.0	90.3	84.4	87.9
IMAZ L	89.7	92.3	81.1	87.7
IMAZ L+B	89.4	90.7	82.3	87.5
THIA L	87.0	88.6	81.0	85.5
THIA L+B	87.7	90.7	80.8	86.4
MEAN	88.6	90.5	81.9	87.0

PLOT AREA HARVESTED 0.00135