

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/W/RN/6 Arable Reference Plots

Rothamsted Research

Rothamsted Research (1981) *80/W/RN/6 Arable Reference Plots* ; Yields Of The Field Experiments 1980, pp 69 - 75 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/W/RN/6

ARABLE REFERENCE PLOTS

Object: To study the long term effects of FYM and N, P and K fertilisers on the yield and mineral content of crops - Woburn Stackyard C.

Sponsor: F.V. Widdowson.

The 21st year, w. barley, w. oats, permanent grass.

For previous years see 60/B/3(t), 61-65/B/2, 66/B/2(t), 67/B/2(t), 68/B/3(t), 69/W/RN/6, 70/W/RN/6(t) and 71-79/W/RN/6.

Design: Blocks of 12 plots for each crop. Barley and oats one third replicate of 3 x 12 x 3 in 3 blocks of 12 plots. Permanent grass 1 block of 12 plots.

Whole plot dimensions: 2.74 x 2.13.

Treatments: All combinations of:-

Blocks

- | | |
|----------|--|
| 1. CROP | Crops in 1980 and previous crops: |
| BW(OW) | W. barley after w. oats in 1979 (arable since 1960) |
| BW(BS) | W. barley after s. barley in 1979 (arable since 1960) |
| BW(BS G) | W. barley after s. barley in 1979 (grass 1960-73, arable thereafter) |
| OW(P) | W. oats after potatoes in 1979 |
| OW(SB) | W. oats after sugar beet in 1979 |
| OW(L) | W. oats after one-year ley in 1979 |

Plots

- | | |
|---------------|---|
| 2. MANURE(79) | Fertilisers and farmyard manure (until 1979, none in 1980): |
| O | |
| N1 | |
| P | |
| N1P | |
| K | |
| NIK | |
| PK | |
| N1PK | |
| N2PK | |
| D | |
| N1PKD | |
| N2PKD | |

N1,2 (kg N): 31.5, 63 (ley): 63, 126 (s. barley and oats): 126, 252 (sugar beet and potatoes): 188, 376 (permanent grass) as ammonium nitrate.

P: P205 at 63 kg as triple superphosphate.

K: K20 at 252 kg as potassium bicarbonate.

D: Farmyard manure at 25 tonnes (permanent grass): 50 tonnes (sugar beet and potatoes): none to other crops.

80/W/RN/6

and some of the combinations of 1 & 2 with:

3. N 80	Nitrogen fertiliser (kg N):
90	25 kg N to seedbed, 30 kg N in February, 35 kg N in April
120	25 kg N to seedbed, 40 kg N in February, 55 kg N in April
150	25 kg N to seedbed, 50 kg N in February, 75 kg N in April

plus one extra block in permanent grass, sown autumn 1973 (PERMGRAS) testing MANURE as above except that treatments were also applied in 1980.

Standard applications:

W. barley and w. oats: Manures: Chalk at 3.7 t. Weedkillers: Ioxynil at 0.32 kg with mecoprop 0.94 kg in 280 l applied twice, with tridemorph on the first occasion, with benomyl on the second. Fungicides: Tridemorph at 0.53 kg. Benomyl at 0.28 kg. Carbendazim at 0.25 kg plus tridemorph at 0.53 kg in 280 l. Insecticide to w. oats only: Phorate at 2.2 kg.

Seed: W. barley: Sonja, sown at 210 kg.

W. oats: Pennal, sown at 210 kg.

Permanent Grass: S215 meadow fescue at 20 kg, S24 perennial ryegrass at 20 kg, crested dogstail at 7 kg, Chewings fescue at 7 kg, smooth stalk meadow fescue at 7 kg, alsike clover at 4 kg, wild white clover at 2 kg. Mixture sown at 67 kg.

Cultivations, etc.:-

W. barley: Rotary cultivated: 3 Sept, 1979. Seed sown, first N applied: 28 Sept. Weedkillers with tridemorph applied, chalk applied: 28 Nov. Second N applied: 18 Feb, 1980. Third N applied, weedkillers with benomyl applied: 1 Apr. Carbendazim plus tridemorph applied: 29 Apr. Harvested: 21 July.

W. oats: Balancing Mg applied and plots dug by hand after potatoes: 2 Oct, 1979. Plots dug by hand after ley: 3 Oct. Balancing Mg applied and plots dug by hand after sugar beet: 17 Oct. Phorate applied, raked in, seed sown, first N applied: 18 Oct. Weedkillers and tridemorph applied, chalk applied: 28 Nov. Second N applied: 18 Feb, 1980. Third N applied, weedkillers with benomyl applied: 1 Apr. Carbendazim plus tridemorph applied: 29 Apr. Harvested: 31 July.

Permanent Grass: P and K applied: 28 Nov, 1979. FYM applied: 11 Mar, 1980. N applied: 11 Mar, 21 May, 8 July. Cut: 21 May, 8 July, 6 Oct.

NOTE: N, P and K contents of grain were measured.

80/W/RN/6

WINTER BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

CROP	BW(OW)	BW(BS)	BW(BS G)	MEAN
MANURE (79)				
O	5.21	3.90	4.50	4.54
N1	3.99	4.08	3.08	3.71
P	5.56	4.49	2.99	4.34
N1P	3.66	3.02	2.56	3.08
K	5.80	4.96	6.83	5.86
N1K	5.97	5.56	5.41	5.65
PK	5.71	5.46	6.63	5.93
N1PK	5.78	5.90	5.27	5.65
N2PK	5.87	5.70	5.89	5.82
D	5.84	6.91	6.58	6.44
N1PKD	6.06	5.80	6.80	6.22
N2PKD	6.35	7.08	6.37	6.60
MEAN	5.48	5.24	5.24	5.32
N 80	90	120	150	MEAN
MANURE (79)				
O	3.90	5.21	4.50	4.54
N1	3.08	4.08	3.99	3.71
P	4.49	5.56	2.99	4.34
N1P	2.56	3.02	3.66	3.08
K	4.96	5.80	6.83	5.86
N1K	5.41	5.56	5.97	5.65
PK	5.46	5.71	6.63	5.93
N1PK	5.27	5.90	5.78	5.65
N2PK	5.87	5.89	5.70	5.82
D	5.84	6.58	6.91	6.44
N1PKD	6.06	6.80	5.80	6.22
N2PKD	6.35	6.37	7.08	6.60
MEAN	4.94	5.54	5.49	5.32

GRAIN MEAN DM% 81.1

80/W/RN/6

WINTER BARLEY

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

CROP MANURE (79)	BW(OW)	BW(BS)	BW(BS G)	MEAN
O	4.77	3.89	4.83	4.50
N1	4.30	3.42	2.95	3.56
P	5.48	4.26	3.43	4.39
N1P	4.30	3.38	3.07	3.59
K	5.32	4.78	6.17	5.42
N1K	5.54	4.88	4.79	5.07
PK	5.52	5.82	6.97	6.11
N1PK	6.21	5.81	5.09	5.70
N2PK	5.81	5.48	7.29	6.19
D	6.00	7.46	8.05	7.17
N1PKD	5.95	6.43	6.58	6.32
N2PKD	6.71	7.01	7.27	7.00
MEAN	5.49	5.22	5.54	5.42
N 80	90	120	150	MEAN
O	3.89	4.77	4.83	4.50
N1	2.95	3.42	4.30	3.56
P	4.26	5.48	3.43	4.39
N1P	3.07	3.38	4.30	3.59
K	4.78	5.32	6.17	5.42
N1K	4.79	4.88	5.54	5.07
PK	5.82	5.52	6.97	6.11
N1PK	5.09	5.81	6.21	5.70
N2PK	5.81	7.29	5.48	6.19
D	6.00	8.05	7.46	7.17
N1PKD	5.95	6.58	6.43	6.32
N2PKD	6.71	7.27	7.01	7.00
MEAN	4.93	5.65	5.68	5.42

STRAW MEAN DM% 57.4

PLOT AREA HARVESTED 0.00050

80/W/RN/6

WINTER OATS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

CROP	OW(P)	OW(SB)	OW(L)	MEAN
MANURE (79)				
O	4.20	4.11	4.96	4.42
N1	3.76	4.12	4.19	4.02
P	4.72	3.77	4.03	4.18
N1P	3.73	3.80	4.20	3.91
K	5.28	4.30	5.91	5.16
N1K	6.22	5.38	4.96	5.52
PK	5.67	5.07	6.12	5.62
N1PK	6.86	6.27	6.20	6.44
N2PK	4.60	6.39	6.09	5.70
D	6.49	6.93	6.98	6.80
N1PKD	6.79	6.26	6.95	6.67
N2PKD	6.57	7.27	6.55	6.79
MEAN	5.41	5.31	5.59	5.44
N 80 90 120 150 MEAN				
MANURE (79)				
O	4.11	4.20	4.96	4.42
N1	4.19	4.12	3.76	4.02
P	3.77	4.72	4.03	4.18
N1P	4.20	3.80	3.73	3.91
K	4.30	5.28	5.91	5.16
N1K	4.96	5.38	6.22	5.52
PK	5.07	5.67	6.12	5.62
N1PK	6.20	6.27	6.86	6.44
N2PK	4.60	6.09	6.39	5.70
D	6.49	6.98	6.93	6.80
N1PKD	6.79	6.95	6.26	6.67
N2PKD	6.57	6.55	7.27	6.79
MEAN	5.10	5.50	5.70	5.44

GRAIN MEAN DM% 72.3

80/W/RN/6

WINTER OATS

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

CROP	OW(P)	OW(SB)	OW(L)	MEAN
MANURE (79)				
O	3.87	3.91	5.13	4.30
N1	3.89	3.88	3.81	3.86
P	4.85	3.83	4.40	4.36
N1P	3.92	4.00	3.59	3.84
K	5.51	4.69	6.56	5.59
N1K	6.29	5.59	5.12	5.67
PK	7.91	6.90	6.86	7.22
N1PK	7.86	7.24	6.81	7.30
N2PK	5.56	6.55	6.78	6.30
D	7.36	7.87	7.26	7.50
N1PKD	9.02	8.24	8.31	8.52
N2PKD	8.74	8.43	8.18	8.45
MEAN	6.23	5.93	6.07	6.08
N 80	90	120	150	MEAN
MANURE (79)				
O	3.91	3.87	5.13	4.30
N1	3.81	3.88	3.89	3.86
P	3.83	4.85	4.40	4.36
N1P	3.59	4.00	3.92	3.84
K	4.69	5.51	6.56	5.59
N1K	5.12	5.59	6.29	5.67
PK	6.90	7.91	6.86	7.22
N1PK	6.81	7.24	7.86	7.30
N2PK	5.56	6.78	6.55	6.30
D	7.36	7.26	7.87	7.50
N1PKD	9.02	8.31	8.24	8.52
N2PKD	8.74	8.18	8.43	8.45
MEAN	5.78	6.12	6.33	6.08

STRAW MEAN DM% 44.0

PLOT AREA HARVESTED 0.00050

80/W/RN/6

PERMANENT GRASS

TONNES/HECTARE

***** TABLES OF MEANS *****

	PERMGRAS : DRY MATTER			
	1ST CUT	2ND CUT	3RD CUT	TOTAL OF 3 CUTS
MANURE (79)				
0	1.65	1.68	1.79	5.12
N1	2.42	2.30	3.24	7.96
P	1.39	1.31	1.39	4.09
N1P	2.66	2.06	2.91	7.62
K	1.73	1.90	2.01	5.64
N1K	2.87	2.94	3.64	9.46
PK	2.05	2.10	2.17	6.33
N1PK	2.78	3.08	3.64	9.50
N2PK	3.83	3.42	4.29	11.54
D	2.93	2.26	2.93	8.13
N1PKD	3.64	3.24	4.32	11.20
N2PKD	4.33	3.89	4.90	13.11
MEAN DM%	22.4	21.2	30.6	24.7