

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1980

[Full Table of Content](#)



80/W/RN/16 Effects of Deep Pk

Rothamsted Research

Rothamsted Research (1981) *80/W/RN/16 Effects of Deep Pk* ; Yields Of The Field Experiments 1980, pp 96 - 97 - DOI: <https://doi.org/10.23637/ERADOC-1-31>

80/W/RN/16

EFFECTS OF DEEP PK

Object: To study the residual effects of subsoiling and of incorporating a large dressing of PK in either the subsoil or topsoil, on yields and nutrient uptakes of s. barley - Woburn Butt Furlong.

Sponsor: J. McEwen.

The seventh year, s. barley.

For previous years see 74-79/W/RN/16.

Design: 4 series of 3 randomised blocks of 4 plots with PREVCROP on series.

Whole plot dimensions: 4.27 x 2.59.

Treatments: All combinations of:-

Series

- | | | |
|-------------|--|--|
| 1. PREVCROP | Previous cropping (1974-1977) (all in barley 1978 & 1979): | |
| P B B | W. wheat, sugar beet, s. barley, potatoes | |
| W B B | Sugar beet, s. barley, potatoes, w. wheat | |
| S B B | S. barley, potatoes, w. wheat, sugar beet | |
| B B B | Potatoes, w. wheat, sugar beet, s. barley | |

Plots

- | | | |
|-----------|---|------------------------------|
| 2. PK SUB | Extra PK and subsoil treatment (applied autumn 1973): | |
| | Extra PK | Subsoil (25-50 cm) treatment |
| - - - | None | None |
| - - S | None | Subsoiled |
| P K T | To topsoil (0-25 cm) | None |
| P K S | To subsoil | Subsoiled |

- NOTES: (1) The rates of P and K were 1930 kg P205, as superphosphate and 460 kg K20 as muriate of potash. These quantities, applied to subsoil, were chosen to equalize available P and K in top and subsoil.
- (2) Subsoiling was done by spade, after removing the topsoil which was then replaced. PK to subsoil was worked in by forking.
- (3) PK to topsoil was applied half before ploughing in autumn half soon after on the plough furrow.

Basal applications: Manures: All Series: Magnesian limestone at 2.5 t; (20:10:10) at 540 kg, combine drilled. Weedkillers: Mecoprop, bromoxynil and ioxynil ('Brittox' 3.5 l in 280 l). Fungicide: Tridemorph ('Beacon' 0.7 l in 280 l).

Seed: Georgie, dressed with ethirimol, sown at 160 kg.

80/W/RN/16

Cultivations, etc.:- Magnesian limestone applied: 31 Oct, 1979. Ploughed: 2 Nov. Spring-tine cultivated with crumbler attached, seed sown: 29 Feb, 1980. Weedkiller applied: 25 Apr. Fungicide applied: 13 May. Harvested by hand: 7 Aug.

NOTE: N, P, K, Na, Ca and Mg contents of grain were measured.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PK SUB PREVCROP	- - -	- - S	P K T	P K S	MEAN
P B B	5.99	6.59	5.78	6.34	6.18
W B B	5.43	5.97	5.31	5.90	5.65
S B B	5.67	6.27	5.45	6.40	5.95
B B B	5.13	5.57	5.69	5.96	5.59
MEAN	5.55	6.10	5.56	6.15	5.84

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PK SUB	PREVCROP* PK SUB
-----	-----	-----
SED	0.173	0.346

* ONLY WHEN COMPARING MEANS WITH SAME LEVELS OF PREVCROP

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	6	0.186	3.2
BLOCK.WP.SP	24	0.423	7.2

GRAIN MEAN DM% 80.7

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

PK SUB PREVCROP	- - -	- - S	P K T	P K S	MEAN
P B B	4.36	4.40	4.07	4.58	4.35
W B B	3.90	4.25	3.54	4.35	4.01
S B B	4.27	4.19	4.03	4.19	4.17
B B B	3.87	3.81	4.65	4.28	4.15
MEAN	4.10	4.16	4.07	4.35	4.17

STRAW MEAN DM% 58.8

SUB PLOT AREA HARVESTED 0.00065