

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/SC/10 and 81/W/CS/10 Long Term Liming - S. Oats

Rothamsted Research

Rothamsted Research (1982) *81/R/SC/10 and 81/W/CS/10 Long Term Liming - S. Oats* ; Yields Of The Field Experiments 1981, pp 102 - 104 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/CS/10 and 81/W/CS/10

LONG TERM LIMING

Object: To study the effects of different amounts of lime on the yields of a sequence of crops. The effects of P & K are also studied - Rothamsted (R) Sawyers I and Woburn (W) Stackyard C.

Sponsors: S. McGrath, D.P. Stribley.

The 20th year, s. oats.

For previous years see 'Details' 1967, 1973 and 74-80/R&W/CS/10.

Design: 2 randomised blocks of 16 plots.

Whole plot dimensions: 6.40 x 18.3.

Treatments: All combinations of:-

1. CHALK Ground chalk (tonnes CaCO₃) (total applied 1962-78):

R	W
0	0
7	6
15	14
30	23

2. P K P & K fertilisers applied (in addition to a basal dressing of 120 kg K in 1981):

(00)0	None
(OK)P1	K annually until 1978, 25 kg P in 1981
(PO)P1	P annually until 1978, 25 kg P in 1981
(PK)P3	P & K annually until 1978, 75 kg P in 1981

NOTES: (1) Until 1978 test P & K were applied cumulatively, rates varied with crop. None was applied in 1979 & 1980 (fallow).
(2) A sub plot test of Mg applied in earlier years has been ignored.

Basal applications:

Sawyers I (R): Manures: N at 80 kg as 'Nitro-Chalk' combine drilled, K at 120 kg as muriate of potash, Mg at 100 kg as kieserite.

Weedkillers: Dicamba, mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l.

Stackyard C (W): Manures: N at 80 kg as 'Nitro-Chalk', K at 120 kg as muriate of potash Mg at 100 kg as kieserite. Weedkillers: Dicamba, mecoprop and MCPA (as 'Banlene Plus' at 4.9 l) in 280 l with the fungicide. Fungicide: Tridemorph at 0.53 l.

Seed: Peniarth, sown at 190 kg (R), 200 kg (W).

81/R/CS/10 and 81/W/CS/10

Cultivations, etc.:-

Sawyers I (R): Basal K and Mg applied: 1 Dec, 1980. P treatments applied, ploughed: 8 Dec. Rotary harrowed, seed sown: 13 Apr, 1981.

Weedkillers applied: 1 June. Combine harvested: 10 Sept.

Stackyard C (W): Ploughed: 13 Nov, 1980, 12 Dec. Basal K and Mg applied: 8 Dec. N applied, heavy spring-tine cultivated: 6 Apr, 1981. Spring-tine cultivated with crumbler attached: 8 Apr. Rotary cultivated, seed sown: 9 Apr. Weedkillers and fungicide applied: 1 June. Combine harvested: 3 Sept.

81/R/CS/10 SAWYERS I(R)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	P K	(00)0	(OK)P1	(PO)P1	(PK)P3	MEAN
CHALK						
0		3.60	2.71	3.70	3.37	3.34
7		3.35	3.67	3.78	3.42	3.55
15		3.32	3.48	3.48	3.86	3.54
30		2.78	3.02	3.26	3.27	3.08
MEAN		3.26	3.22	3.55	3.48	3.38

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHALK	P K	CHALK P K
SED	0.139	0.139	0.278

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	0.278	8.2

GRAIN MEAN DM% 82.3

PLOT AREA HARVESTED 0.00520

81/W/CS/10 STACKYARD C (W)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	P K	(00)0	(0K)P1	(P0)P1	(PK)P3	MEAN
CHALK						
0		3.56	3.90	4.05	4.15	3.92
6		3.60	3.65	3.99	3.95	3.80
14		3.95	3.45	3.35	4.04	3.70
23		3.61	3.66	3.62	3.52	3.60
MEAN		3.68	3.66	3.75	3.92	3.75

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHALK	P K	CHALK P K

SED	0.124	0.124	0.248

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	0.248	6.6
GRAIN MEAN DM%	82.5		
PLOT AREA HARVESTED	0.00503		