

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/CS/165 Sclerotinia Control - Red and White Clover

Rothamsted Research

Rothamsted Research (1982) *81/R/CS/165 Sclerotinia Control - Red and White Clover* ; Yields Of The Field Experiments 1981, pp 136 - 137 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/CS/165

SCLEROTINIA CONTROL

Object: To study the effects of two fungicides and paraquat on the incidence of Sclerotinia and on yield of red and white clover - Little Knott I.

Sponsor: J.F. Jenkyn.

The fourth year of treatments, red and white clover.

For previous years see 78-80/R/CS/165.

Design: 2 randomised blocks of 30 plots.

Whole plot dimensions: 2.13 x 3.05.

Treatments: All combinations of:-

1. VARIETY Varieties and their resistance to Sclerotinia trifoliorum:

 BLANC WR Blanca, white clover, resistant
 SABED WS Sabeda, white clover, susceptible
 HUNGA RR Hungaropoly, red clover, resistant
 SABTO RS Sabtoron, red clover, susceptible

2. TREATMNT Chemical sprays and timing (cumulative):

 NONE None
 IPROD E Iprodione early period, 26 Sept, 1980 and 4 Nov.
 IPROD M Iprodione mid period, 4 Nov and 27 Nov.
 IPROD L Iprodione late period, 27 Nov and 18 Dec.
 IPROD A Iprodione all periods, 26 Sept, 4 Nov, 27 Nov, 18 Dec, 1980, 26 Jan, 1981
 BENOMY A Benomyl all periods, 26 Sept, 4 Nov, 27 Nov, 18 Dec, 1980, 26 Jan, 1981
 PARAQ W Paraquat, 27 Nov, 1979 and 2 Apr, 1981

plus two extra treatments (cumulative):

EXTRA

- | | |
|----------|---|
| SABED PS | Sabeda, sprayed paraquat in spring, 2 Apr, 1981 |
| SABTO PS | Sabtoron, sprayed paraquat in spring, 2 Apr |

- NOTES: (1) Chemical treatments were applied, in 340 l, as follows:-
Iprodione at 0.50 kg, benomyl at 0.50 kg (both applied with 'Spreadite', a wetting agent, at 0.25 kg). Paraquat at 0.28 kg ion.
- (2) On 27 Nov one of the replicates of the treatment combinations of HUNGA RR with NONE and IPROD M received paraquat in error and one of the replicates of the treatment combinations of HUNGA RR and SABTO RS with PARAQ W did not receive paraquat by the same error. The omitted treatment was applied on 8 Dec.
- (3) The proportion of weeds to clover was assessed.

81/R/CS/165

Basal applications: Manures: (0:14:28) at 540 kg. Weedkiller: Propyzamide at 0.7 kg in 340 l.

Cultivations, etc.: - Weedkiller applied: 26 Sept, 1980. PK applied: 4 Dec. Cut: 6 July, 1981.

1ST CUT (6/7/81) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

VARIETY TREATMNT	BLANC WR	SABED WS	HUNGA RR	SABTO RS	MEAN
NONE	3.90	3.62	5.23	5.32	4.52
IPROD E	3.64	3.38	6.18	4.52	4.43
IPROD M	3.73	4.11	5.00	5.59	4.61
IPROD L	3.31	3.45	5.60	4.51	4.22
IPROD A	3.35	3.94	5.35	6.08	4.68
BENOMY A	4.55	4.67	5.32	6.02	5.14
PARQ W	3.60	4.21	5.07	4.54	4.35
MEAN	3.72	3.91	5.39	5.23	4.56

EXTRA	SABED PS	SABTO PS	MEAN
	3.49	3.87	3.68

GRAND MEAN 4.50

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	TREATMNT	VARIETY	TREATMNT VARIETY & EXTRA

SED	0.735	0.368	0.278	0.735

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	29	0.735	16.3

1ST CUT MEAN DM% 22.2

PLOT AREA HARVESTED 0.00028