

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1981

[Full Table of Content](#)



## 81/R/CS/203 Species Mixtures and Phialophora - W. Wheat

### Rothamsted Research

Rothamsted Research (1982) *81/R/CS/203 Species Mixtures and Phialophora - W. Wheat* ; Yields Of The Field Experiments 1981, pp 163 - 164 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/CS/203

SPECIES MIXTURES AND PHIALOPHORA

Object: To study the effects of lucerne, grass and clover, singly and in mixtures, and of soil inoculation on populations of *Phialophora radicularis* and on take-all (*Gaeumannomyces graminis*) and yield of following w. wheat crops - Stubbings.

Sponsors: E. Lester, D.B. Slope, R.J. Gutteridge.

The fifth year, w. wheat.

For previous years see 79-80/R/CS/203.

Design: 4 randomised blocks of 9 plots, split into 4.

Whole plot dimensions: 4.27 x 27.1.

Treatments: All combinations of:-

1. Whole plots

CRP INOC	Crops in 1977 and 1978 (all w. wheat 1979 to 1981) and inoculation:-	
	1977	1978
C C	White clover	White clover
G G	Ryegrass	Ryegrass
GC GC	Ryegrass/white clover mixture	Ryegrass/white clover
LU LU	Lucerne	Lucerne
LU LU I	Lucerne	Lucerne + Prg inoculum to 1979 wheat
GLU GLU	Ryegrass/lucerne in alternate rows	Ryegrass/lucerne
W G	Spring wheat	Ryegrass, sown into spring wheat stubble
WG G	Spring wheat undersown with ryegrass	Ryegrass
WGI G	Spring wheat, inoculated Prg, undersown with ryegrass	Ryegrass

2. Sub plots

N	Nitrogen fertiliser (kg N) from 1979 to 1981, cumulative:
0	
50	
100	
150	

NOTE: The inoculum used for the I treatments was an agar culture of Prg mixed with sand. It was broadcast and power harrowed into the soil before sowing.

81/R/CS/203

Basal applications: Manures: (0:20:20) at 250 kg, combine drilled.  
 Weedkillers: Chlortoluron at 5.6 l in 250 l. Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 3.7 l) in 250 l. Fungicide: Prochloraz at 0.4 l in 250 l.

Seed: Flanders, sown at 200 kg.

Cultivations, etc.: - Ploughed: 29 Aug, 1980. Spring-tine cultivated: 30 Sept. Rotary harrowed: 1 Oct. Seed sown: 2 Oct. Chlortoluron applied: 3 Oct. N and 'Brittox' applied: 17 Apr, 1981. Fungicide applied: 3 June. Combine harvested: 25 Aug.

NOTE: Take-all was assessed in April and July. Bio-assays of soils for take-all were made after harvest.

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	N	0	50	100	150	MEAN
CRP INOC						
C C		1.87	2.93	3.82	3.86	3.12
G G		3.16	4.97	4.86	4.64	4.41
GC GC		2.86	5.06	5.61	5.16	4.67
LU LU		1.94	3.06	3.20	3.12	2.83
LU LU I		1.83	2.89	3.50	3.74	2.99
GLU GLU		2.91	3.73	4.65	4.24	3.88
W G		2.83	4.15	3.92	4.41	3.83
WG G		2.90	4.43	4.70	4.64	4.17
WGI G		2.86	4.04	4.98	4.65	4.13
MEAN		2.57	3.92	4.36	4.27	3.78

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	CRP INOC	N	CRP INOC
			N
SED	0.246	0.136	0.432
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
CRP INOC			0.409

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	24	0.349	9.2
BLOCK.WP.SP	81	0.579	15.3

GRAIN MEAN DM% 82.2

SUB PLOT AREA HARVESTED 0.00127