

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/CS/250 Control of Cephalosporium - W. Wheat

Rothamsted Research

Rothamsted Research (1982) *81/R/CS/250 Control of Cephalosporium - W. Wheat* ; Yields Of The Field Experiments 1981, pp 191 - 192 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/CS/250

CONTROL OF CEPHALOSPORIUM

Object: To study the effects of aldicarb, benomyl and HCH on the incidence of Cephalosporium Stripe (*Cephalosporium gramineum*) and on the yield of w. wheat - New Zealand.

Sponsors: R.J. Gutteridge, K.E. Fletcher.

The second year, w. wheat.

For previous year see 80/R/CS/250.

Design: 3 randomised blocks of 8 plots.

Whole plot dimensions: 2.67 x 6.10.

Treatments: All combinations of:-

1. CHEMICAL Chemicals and methods of application:
 ALDICARB Aldicarb at 10 kg worked in to seedbed
 BENOMYL Benomyl at 20 kg worked in to seedbed
 HCH HCH seed dressing at 0.4 g per kg seed
2. YEARS Harvest years of application:
 1980+81 1980 and 1981 cumulative treatment
 1981 1981, none in 1980

plus one extra treatment:

EXTRA

NONE No chemical treatment in 1980 & 1981 (duplicated)

NOTES: (1) 1981 soil treatments were applied on 2 Oct, 1980.
(2) 1981 benomyl was applied as a drench in 11,200 l of water.

Basal applications: Manures: (10:23:23) at 250 kg, combine drilled.
'Nitro-Chalk' at 390 kg. Weedkillers: Paraquat at 0.84 kg ion in 250 l.
Chlortoluron at 5.6 l in 250 l. Fungicide: Prochloraz at 0.4 l in 250 l.
Growth regulator: Chlormequat at 1.7 l in 250 l.

Seed: Flanders, sown at 200 kg.

Cultivations, etc.: - Paraquat applied: 19 Sept, 1980. Ploughed: 27 Sept.
Spring-tine cultivated twice: 1 Oct. Treatments applied, spike rotary cultivated: 2 Oct. Seed sown: 3 Oct. Chlortoluron applied: 10 Oct. N applied: 16 Apr, 1981. Growth regulator applied: 24 Apr. Fungicide applied: 5 May. Combine harvested: 26 Aug.

NOTE: Incidence of Cephalosporium, take-all and Phialophora was measured in June. In April soil samples were examined for wireworm larvae, and plants for wireworm damage.

81/R/CS/250

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

CHEMICAL YEARS	ALDICARB	BENOMYL	HCH	MEAN
1980+81	7.53	7.47	8.00	7.67
1981	7.86	8.22	7.72	7.93
MEAN	7.69	7.84	7.86	7.80

NONE 7.75

GRAND MEAN 7.79

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	YEARS	CHEMICAL	YEARS CHEMICAL

SED	0.244	0.298	0.422

SED FOR NONE AGAINST ANY ITEM IN CHEMICAL.YEARS TABLE IS 0.365

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	0.517	6.6

GRAIN MEAN DM% 83.4

PLOT AREA HARVESTED 0.00161