

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/CS/261 Benomyl and Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (1982) *81/R/CS/261 Benomyl and Take-all - W. Wheat* ; Yields Of The Field Experiments 1981, pp 204 - 205 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/CS/261

BENOMYL AND TAKE-ALL

Object: To study the effects of benomyl, applied to the soil with and without a surfactant, and of nuarimol on take-all (*Gaeumannomyces graminis*) and on the yield of w. wheat - New Zealand.

Sponsor: G.L. Bateman.

The first year, w. wheat.

Design: 4 randomised blocks of 7 plots.

Whole plot dimensions: 2.13 x 12.2.

Treatments:

FUNGICIDE	Fungicides, surfactant and times and methods of application:
NONE	None
SF S	Surfactant (an alcohol ethoxylate at 115 l) as a drench in 11,500 l in spring
BEN A	Benomyl at 20 kg worked in to seedbed in autumn
BEN S	Benomyl at 20 kg, as a drench in 11,500 l in spring
BEN+SF S	Benomyl at 20 kg plus surfactant as above as a drench in 11,500 l in spring
NUA A	Nuarimol at 2.2 kg worked in to seedbed in autumn
NUA S	Nuarimol at 2.2 kg, as a drench in 11,500 l in spring

NOTE: Treatments were applied on 2 Oct, 1980 and 24 Apr, 1981.

Basal applications: Manures: (10:23:23) at 250 kg, combine drilled. 'Nitro-Chalk' at 390 kg. Weedkillers: Paraquat at 0.84 kg ion in 250 l. Chlortoluron at 5.6 l in 250 l. Growth regulator: Chlormequat at 1.7 l in 250 l.

Seed: Flanders, with no seed dressing, sown at 200 kg.

Cultivations, etc.: - Paraquat applied: 19 Sept, 1980. Ploughed: 27 Sept. Spring-tine cultivated twice: 1 Oct. Spike rotary cultivated, harrowed: 2 Oct. Seed sown: 3 Oct. Chlortoluron applied: 10 Oct. N applied: 16 Apr, 1981. Growth regulator applied: 24 Apr. Combine harvested: 25 Aug. Previous crops: Grass and fallow 1979, w. wheat, 1980.

NOTE: Foot and root rots were assessed monthly between April and July.

81/R/CS/261

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE	NONE	SF S	BEN A	BEN S	BEN+SF S	NUA A	NUA S	MEAN
	7.64	5.02	7.48	8.59	6.05	7.70	7.72	7.17

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGCIDE
-----	-----
SED	0.367

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	18	0.520	7.2

GRAIN MEAN DM% 79.2

PLOT AREA HARVESTED 0.00176