

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1981

[Full Table of Content](#)



## 81/R/WW/2 Seed Rates & Divided N Dressings - W. Wheat

### Rothamsted Research

Rothamsted Research (1982) *81/R/WW/2 Seed Rates & Divided N Dressings - W. Wheat* ; Yields Of The Field Experiments 1981, pp 226 - 228 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/WW/2

WINTER WHEAT

SEED RATES & DIVIDED N DRESSINGS

Object: To study the effects of a range of rates of early nitrogen dressings on the growth and yield of wheat sown at half or standard seed rate - Gt. Harpenden I.

Sponsors: J. McEwen, R. Moffitt.

Design: 2 randomised blocks of 30 plots.

Whole plot dimensions: 4.27 x 8.08.

Treatments: All combinations of:-

1. SD RATE      Seed rates (kg):

100  
200

2. EARLY N      Nitrogen fertiliser applied 16 Feb, 1981 (kg N):

0  
25  
50  
75

3. APRIL N      Nitrogen fertiliser applied 3 Apr, 1981 (kg N):

75  
100  
125

plus extra treatments, all combinations of:-

1. SD RATEX      Seed rates (kg):

100  
200

2. APRIL NX      Nitrogen fertiliser applied 3 Apr, 1981 (kg N):

150  
175  
200

Basal applications: Manures: (0:20:20) at 310 kg, combine drilled.  
Weedkillers: Chlortoluron at 5.6 l in 250 l. Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 3.7 l) in 250 l. Fungicides: Prochloraz at 0.4 l in 250 l applied twice, with the growth regulator on the first occasion and with maneb and zineb on the second. Maneb at 1.2 kg with zineb at 0.13 kg. Growth regulator: Chlormequat at 1.7 kg.

Seed: Flanders.

81/R/WW/2

Cultivations, etc.:— Disc harrowed twice: 12 Sept, 1980. Heavy spring-tine cultivated three times: 6 Oct. Spring-tine cultivated: 8 Oct. Seed sown: 9 Oct. Chlortoluron applied: 14 Oct. 'Brittox' applied: 21 Apr, 1981. Growth regulator and prochloraz applied: 5 May. Prochloraz, maneb and zineb applied: 19 June. Combine harvested: 25 Aug. Previous crops: S. barley, 1979, w. oats, 1980.

NOTES: (1) Plant counts were made in November, shoot counts in February and April and ear counts in July.  
 (2) 1000 grain weights and N content of grain were measured.

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

EARLY N	0	25	50	75	MEAN
SD RATE					
100	7.94	8.02	8.18	8.38	8.13
200	7.82	8.24	8.18	8.41	8.16
MEAN	7.88	8.13	8.18	8.39	8.15
APRIL N	75	100	125	MEAN	
SD RATE					
100	7.93	8.27	8.18	8.13	
200	7.97	8.28	8.25	8.16	
MEAN	7.95	8.27	8.22	8.15	
APRIL N	75	100	125	MEAN	
EARLY N					
0	7.57	8.08	8.00	7.88	
25	7.86	8.20	8.32	8.13	
50	7.94	8.27	8.33	8.18	
75	8.42	8.54	8.22	8.39	
MEAN	7.95	8.27	8.22	8.15	
SD RATE	APRIL N	75	100	125	
100	EARLY N				
	0	7.71	8.00	8.13	
	25	7.76	8.13	8.16	
	50	7.83	8.43	8.28	
	75	8.43	8.53	8.17	
200	0	7.44	8.17	7.87	
	25	7.96	8.27	8.48	
	50	8.06	8.11	8.38	
	75	8.41	8.56	8.27	
APRIL NX	150	175	200	MEAN	
SD RATEX					
100	8.19	7.94	8.22	8.11	
200	8.08	8.09	8.23	8.13	
MEAN	8.13	8.01	8.22	8.12	
GRAND MEAN	8.14				

81/R/WW/2

GRAIN TONNES/HECTARE

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	SD RATE	EARLY N	APRIL N	SD RATEX
SED	0.075	0.106	0.092	0.150

TABLE	APRIL NX	SD RATE EARLY N	SD RATE APRIL N	EARLY N APRIL N
SED	0.184	0.150	0.130	0.184

TABLE	SD RATE EARLY N APRIL N	SD RATEX APRIL NX
	0.260	0.260

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	29	0.260	3.2

GRAIN MEAN DM% 84.2

PLOT AREA HARVESTED 0.00246