

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1981

[Full Table of Content](#)



## 81/R/WW/4 Fungicides and Microflora - W. Wheat

### Rothamsted Research

Rothamsted Research (1982) *81/R/WW/4 Fungicides and Microflora - W. Wheat* ; Yields Of The Field Experiments 1981, pp 250 - 252 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/WW/4

WINTER WHEAT

FUNGICIDES AND MICROFLORA

Object: To study the effects of a range of fungicides and times of application on the colonisation of fungi on flag leaves and developing ears and on the yield of w. wheat - Webbs.

Sponsor: N. Magan.

Design: 2 randomised blocks of 24 plots.

Whole plot dimensions: 4.27 x 16.2.

Treatments: All combinations of:-

1. E FUNG Early-applied fungicides:

NONE	None
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg applied on 1 June, 1981

2. L FUNG A Late-applied fungicides:

BENOMYL	Benomy1 at 0.56 kg
CAPTAFOL	Captafol at 1.4 kg
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg
IMAZALIL	Imazalil at 0.39 kg
PROCHLOR	Prochloraz at 0.40 l

3. LFNGDATE Dates of applying late fungicide:

12 JUNE	12 June, 1981
22 JUNE	22 June

plus two extra treatments not given L FUNG:

L FUNG 0

NONE	No early-applied fungicide (duplicated)
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg applied on 1 June, 1981 (duplicated)

NOTE: Treatment sprays were applied in 340 l.

Basal applications: Manures: (0:20:20) at 310 kg, combine drilled. 'Nitro-Chalk' at 390 kg. Weedkillers: Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 3.7 l) in 250 l.

Seed: Maris Huntsman, sown at 200 kg.

Cultivations, etc.: - Heavy spring-tine cultivated twice: 2 Oct, 1980. Spring-tine cultivated, seed sown: 3 Oct. N applied: 8 Apr, 1981. Weedkillers applied: 10 Apr. Combine harvested: 29 Aug. Previous crops: W. wheat 1979, potatoes, 1980.

81/R/WW/4

NOTE: Grain and flag leaf microflora, especially *Alternaria* and *Cladosporium*, were assessed at fortnightly intervals after heading. Thousand grain weights were measured, and grain was assessed for germination and seedling growth.

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

LFNGDATE	12 JUNE	22 JUNE				MEAN
E FUNG						
NONE	8.21	8.31				8.26
CARB+MAN	8.45	8.41				8.43
MEAN	8.33	8.36				8.34
L FUNG A	BENOMYL	CAPTAFOL	CARB+MAN	IMAZALIL	PROCHLOR	MEAN
E FUNG						
NONE	8.39	8.27	8.25	7.99	8.38	8.26
CARB+MAN	8.40	8.32	8.59	8.24	8.59	8.43
MEAN	8.40	8.29	8.42	8.11	8.49	8.34
L FUNG A	BENOMYL	CAPTAFOL	CARB+MAN	IMAZALIL	PROCHLOR	MEAN
LFNGDATE						
12 JUNE	8.47	8.22	8.28	8.11	8.56	8.33
22 JUNE	8.32	8.36	8.56	8.12	8.41	8.36
MEAN	8.40	8.29	8.42	8.11	8.49	8.34
E FUNG	L FUNG A	BENOMYL	CAPTAFOL	CARB+MAN	IMAZALIL	PROCHLOR
NONE	LFNGDATE					
	12 JUNE	8.49	8.21	8.13	7.89	8.31
	22 JUNE	8.30	8.33	8.37	8.08	8.46
CARB+MAN	12 JUNE	8.46	8.23	8.43	8.32	8.81
	22 JUNE	8.34	8.40	8.75	8.17	8.37
L FUNG 0	NONE	CARB+MAN				
	7.93	8.00				

GRAND MEAN 8.28

81/R/WW/4

GRAIN TONNES/HECTARE

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	L FUNG O	E FUNG	LFNGDATE	L FUNG A
SED	0.172	0.077	0.077	0.121
TABLE	E FUNG LFNGDATE	E FUNG L FUNG A	LFNGDATE L FUNG A	E FUNG LFNGDATE L FUNG A
SED	0.109	0.172	0.172	0.243

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	25	0.243	2.9

GRAIN MEAN DM% 83.7

SUB PLOT AREA HARVESTED 0.00260