

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)

International Agricultural Institute
Rothamsted
Harpenden
Herts
SG8 5LR
UK
Tel: +44 (0)1494 473900
Fax: +44 (0)1494 473901
Email: eradoc@rothamsted.ac.uk
Web: <http://www.rothamsted.ac.uk/eradoc/>

81/R/WW/9 Nitrification Inhibitors and Soil N - W. Wheat

Rothamsted Research

Rothamsted Research (1982) *81/R/WW/9 Nitrification Inhibitors and Soil N - W. Wheat* ; Yields Of The Field Experiments 1981, pp 261 - 262 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/WW/9

WINTER WHEAT

NITRIFICATION INHIBITORS AND SOIL N

Object: To study the effects of nitrification inhibitors, applied before sowing on the leaching of nitrogen in the soil and on the yield of w. wheat - Gt. Harpenden I.

Sponsor: G.A. Rodgers.

Design: 2 randomised blocks of 21 plots.

Whole plot dimensions: 2.74 x 11.6.

Treatments: All combinations of:-

1. N INHIB Nitrification inhibitors before sowing:

NONE	None
DICYAN 1	Dicyandiamide at 5 kg
DICYAN 2	Dicyandiamide at 20 kg
ETRIDI 1	Etridiazole at 0.5 kg
ETRIDI 2	Etridiazole at 2.0 kg
NITRAP 1	Nitrapyrin at 0.5 kg
NITRAP 2	Nitrapyrin at 2.0 kg

2. SPRING N Nitrogen fertiliser (kg N) in spring on 16 April, 1981:

0
35
70

NOTE: Nitrification inhibitors were applied on 24 Sept, 1980.

Basal applications: Manures: (0:20:20) at 310 kg, combine drilled.

Weedkillers: Isoproturon at 2.5 kg in 250 l. Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 3.7 l) in 250 l. Fungicides: Prochloraz at 0.4 l applied with maneb at 1.2 kg and zineb at 0.13 kg in 250 l.

Seed: Flanders, sown at 190 kg.

Cultivations, etc.: - Subsoiled four times: 13 June, 1980. Chisel ploughed twice: 16 June. Rotary cultivated: 18 June, 8 July, 7 Aug, 18 Sept. Rotary harrowed: 24 Sept. Seed sown: 29 Sept. Isoproturon applied: 4 Oct. 'Brittox' applied: 21 Apr, 1981. Fungicides applied: 22 June. Combine harvested: 26 Aug. Previous crops: Grass 1979 and 1980.

NOTES: (1) Soil cores were taken to 20 cms, at 21 day intervals between September and April, and to 1 m in January, and analysed for ammonium and nitrate concentrations.

(2) Dicyandiamide concentrations were measured on several occasions after applications. Total N was measured in plants in April, and ears were counted in June. Harvested grain and straw was analysed for total N.

81/R/WW/9

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

SPRING N N INHIB	0	35	70	MEAN
NONE	4.55	5.87	7.10	5.84
DICYAN 1	4.69	6.62	7.21	6.17
DICYAN 2	5.30	6.40	7.18	6.29
ETRIDI 1	4.36	6.13	7.46	5.98
ETRIDI 2	5.08	5.84	7.14	6.02
NITRAP 1	5.10	6.40	6.48	6.00
NITRAP 2	4.88	6.17	6.75	5.93
MEAN	4.85	6.20	7.05	6.03

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	N INHIB	SPRING N	N INHIB SPRING N
-----	-----	-----	-----
SED	0.282	0.185	0.488

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	20	0.488	8.1

GRAIN MEAN DM% 86.0

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

SPRING N N INHIB	0	35	70	MEAN
NONE	2.32	3.19	3.49	3.00
DICYAN 1	2.34	2.81	3.49	2.88
DICYAN 2	2.49	3.20	3.67	3.12
ETRIDI 1	2.36	3.07	3.99	3.14
ETRIDI 2	2.80	2.67	3.83	3.10
NITRAP 1	2.36	3.31	3.51	3.06
NITRAP 2	2.28	3.34	3.29	2.97
MEAN	2.42	3.09	3.61	3.04

STRAW MEAN DM% 88.2

PLOT AREA HARVESTED 0.00156