

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1981

[Full Table of Content](#)



## 81/W/B/1 Mildew Study - W. & S. Barley

### Rothamsted Research

Rothamsted Research (1982) *81/W/B/1 Mildew Study - W. & S. Barley ; Yields Of The Field Experiments 1981*, pp 273 - 276 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/W/B/1

WINTER & SPRING BARLEY

MILDEW STUDY

Object: To study the effects of fungicides applied to w. & s. barley on the incidence of mildew and on yield and whether these effects are influenced by neighbouring treatments - Woburn Far Field II.

Sponsor: D.W. Hollomon.

Design: W. barley: 3 blocks of 7 plots split into 2  
S. barley: 3 blocks of 7 plots

Whole plot dimensions: 8.53 x 8.53.

Treatments:

To W. BARLEY: All combinations of:-

Whole plots

1. SEEDRESS      Seed dressing to w. barley and one adjacent plot of s. barley, other adjacent 'plot' of s. barley sprayed tridemorph at 0.53 kg in 280 l on 30 May:

WBO SBO	None to w. and s. barley
WBO SBE	None to w. barley, ethirimol to s. barley
WBO SBT	None to w. barley, triadimenol + fuberidazole to s. barley
WBE SBO	Ethirimol to w. barley, none to s. barley
WBE SBE	Ethirimol to w. and s. barley
WBT SBO	Triadimenol + fuberidazole to w. barley, none to s. barley
WBT SBT	Triadimenol + fuberidazole to w. and s. barley

Sub plots

2. POSITION      Position of w. barley plots in relation to s. barley plots testing seed dressing:

N EAST	North east
S WEST	South west

To S. BARLEY:

SEEDRESS      Seed dressings to s. barley and to both adjacent plots of w. barley:

SBO WBO	None to s. and w. barley
SBO WBE	None to s. barley, ethirimol to w. barley
SBO WBT	None to s. barley, triadimenol + fuberidazole to w. barley
SBE WBO	Ethirimol to s. barley, none to w. barley
SBE WBE	Ethirimol to s. and w. barley
SBT WBO	Triadimenol + fuberidazole to s. barley, none to w. barley
SBT WBT	Triadimenol + fuberidazole to s. and w. barley

81/W/B/1

Standard applications: Manures: (10:23:23) at 250 kg for w. barley, (0:20:20) at 310 kg for s. barley, N at 130 kg as 'Nitro-Chalk' to all plots. Weedkillers: Mecoprop with bromoxynil and ioxynil (as 'Brittox' at 3.5 l) in 280 l to w. barley, isoproturon at 2.5 kg in 280 l to w. barley, dicamba with mecoprop and MCPA (as 'Banlene Plus' at 4.9 l) in 280 l to s. barley.

Seed: W. barley: Hoppel, sown at 160 kg.  
S. barley: Wing, sown at 160 kg.

Cultivations, etc.: - Deep-tine cultivated: 24 Sept, 1980. NPK applied and spring-tine cultivated with crumbler attached for w. barley: 1 Oct. W. barley seed sown: 2 Oct. Deep-tine cultivated for s. barley: 31 Oct. PK applied for s. barley: 26 Feb, 1981. Spring-tine cultivated with crumbler attached three times for s. barley: 27 Feb, 6 Apr, 7 Apr. N applied to all plots: 20 Mar. 'Brittox' applied to w. barley: 27 Mar. S. barley, seed sown: 8 Apr. Isoproturon applied to w. barley: 22 Apr. 'Banlene Plus' applied to s. barley: 29 May. W. barley combine harvested: 28 July. S. barley combine harvested: 18 Aug. Previous crops: Grain legumes 1979, w. wheat 1980.

NOTE: The incidence of mildew (*Erysiphe graminis*) was measured during December and April for w. barley and on four occasions between May and July for s. barley.

81/W/B/1

WINTER BARLEY

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

POSITION SEEDRESS	N EAST	S WEST	MEAN
WBO SBO	6.41	6.56	6.49
WBO SBE	7.18	6.72	6.95
WBO SBT	7.22	7.24	7.23
WBE SBO	6.72	7.50	7.11
WBE SBE	6.88	7.11	6.99
WBT SBO	6.94	7.05	6.99
WBT SBT	6.53	6.76	6.64
MEAN	6.84	6.99	6.91

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	SEEDRESS	POSITION	SEEDRESS POSITION
SED	0.303	0.100	0.356
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
SEEDRESS			0.265

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.371	5.4
BLOCK.WP.SP	14	0.324	4.7

GRAIN MEAN DM% 79.6

SUB PLOT AREA HARVESTED 0.00243

81/W/B/1

SPRING BARLEY

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

SEEDRESS	
SBO WBO	5.78
SBO WBE	6.07
SBO WBT	5.78
SBE WBO	6.10
SBE WBE	6.00
SBT WBO	6.13
SBT WBT	6.16
MEAN	6.00

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	SEEDRESS
SED	0.178

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.217	3.6

GRAIN MEAN DM% 87.6

PLOT AREA HARVESTED 0.00243