

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/BE/6 Effects of Pests and Pathogens - S. Beans

Rothamsted Research

Rothamsted Research (1982) *81/R/BE/6 Effects of Pests and Pathogens - S. Beans* ; Yields Of The Field Experiments 1981, pp 302 - 303 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/BE/6

SPRING BEANS

EFFECTS OF PESTS AND PATHOGENS

Object: To assess the benefits from three amounts of pest and disease control on irrigated and unirrigated s. beans - Fosters Corner.

Sponsors: J. McEwen, R. Bardner, A.J. Cockbain, J.M. Day, W. Day, K.E. Fletcher, G.A. Salt, R.M. Webb, J.F. Witty, D.P. Yeoman.

Design: 4 randomised blocks of 2 plots split into 3.

Whole plot dimensions: 4.27 x 13.7.

Treatments: all combinations of:-

Whole plots

1. IRRIGATN Irrigation:
 NONE None
 FULL Full (total 49 mm)

Sub plots

2. PATHCONT Pest and pathogen control (in addition to basal):
 STANDARD None
 ENHANCED Phorate at 2.24 kg, combine drilled
 Benomy1 at 0.56 kg on 11 Aug
 FULL Aldicarb at 10 kg on 19 Feb, 1981
 Fosetyl-A1 at 2.24 kg on 23 Apr
 Benomy1 at 0.56 kg on 28 July
 Benomy1 at 0.56 kg on 11 Aug

NOTES: (1) Treatment sprays were applied in 340 l.
(2) Irrigation was applied to reduce a deficit of 50 mm to 25 mm before pod set, and from 80 mm to 55 mm after pod set. (mm water):

3 July	12
4 July	12
10 July	<u>25</u>
Total	49

Basal applications: Manures: Chalk at 7.5 t. Weedkiller: Simazine at 1.2 kg in 250 l. Insecticide: Pirimicarb at 0.14 kg in 250 l applied twice.

Seed: Minden, sown at 240 kg (500,000 seeds per hectare).

Cultivations, etc.: - Chalk applied: 3 Dec, 1980. Ploughed: 10 Dec. Spring-tine cultivated three times: 18 Feb, 1981, 19 Feb. Rotary harrowed, seed sown: 20 Feb. Weedkiller applied: 8 Apr. Basal insecticide applied: 3 June, 28 July. Combine harvested: 8 Sept. Previous crops: W. wheat 1979, s. barley 1980.

81/R/BE/6

NOTE: Plant counts were made after establishment and components of yield measured before harvest. Total above-ground dry matter and N content were measured in August. Migratory nematodes, root and foliar fungi, aphids, weevils and viruses were counted at intervals during the season. N content of grain was measured.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PATHCONT IRRIGATN	STANDARD	ENHANCED	FULL	MEAN
NONE	3.93	4.50	4.65	4.36
FULL	4.66	5.00	5.02	4.89
MEAN	4.29	4.75	4.84	4.63

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PATHCONT	IRRIGATN* PATHCONT
SED	0.074	0.104

* WITHIN THE SAME LEVEL OF IRRIGATN ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	12	0.147	3.2

GRAIN MEAN DM% 86.8

SUB PLOT AREA HARVESTED 0.00293