Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1981



Full Table of Content

81/R/BE/19 Electrostatic Spraying and Aphids - S. Beans

Rothamsted Research

Rothamsted Research (1982) 81/R/BE/19 Electrostatic Spraying and Aphids - S. Beans; Yields Of The Field Experiments 1981, pp 321 - 322 - DOI: https://doi.org/10.23637/ERADOC-1-35

81/R/BE/19

SPRING BEANS

ELECTROSTATIC SPRAYING AND APHIDS

Object: To study the effects of applying aphicides with an electrostatic sprayer on the control of black fly (Aphis fabae) and on the yield of s.beans - Whittlocks.

Sponsors: D.C. Griffiths, A.J. Arnold, G.R.Cayley, P. Etheridge, F.T. Phillips, B. Pye, G.C. Scott.

Design: 4 randomised blocks of 10 plots.

Whole plot dimensions: 3.20 x 9.14.

Treatments:

TREATMNT	Sprayers and aphicides:
NONE	None
EL V2 D1	Electrostatic sprayer, droplets charged 25 kV, demeton-s-methyl at 0.06 kg.
EL V2 D2	Electrostatic sprayer, droplets charged 25 kV, demeton-s-methyl at 0.12 kg.
EL V 2 D4	Electrostatic sprayer, droplets charged 25 kV, demeton-s-methyl at 0.24 kg.
EL V2 P4	Electrostatic sprayer, droplets charged 25 kV, permethrin at 0.10 kg.
EL - U	Electrostatic sprayer, droplets uncharged, 'Uvitex' dye
EL V1 U	Electrostatic sprayer, droplets charged 15 kV, 'Uvitex' dye
EL V2 U	Electrostatic sprayer, droplets charged 25 kV, 'Uvitex' dye
ST - D4	Standard sprayer, droplets uncharged, demeton-s-methyl at 0.24 kg.
ST - P4	Standard sprayer, droplets uncharged, permethrin at 0.10 kg.

NOTE: Treatments were applied on 12 June, 1981 in 3.73 l of oil or water by electrostatic sprayer, in 450 l of water by standard sprayer.

Basal applications: Manure: Chalk at 7.5 t. Weedkillers: Simazine at 1.2 kg in 250 l. Benomyl at 0.55 kg in 250 l.

Seed: Minden, sown at 260 kg.

Cultivations, etc.:- Chalk applied: 2 Sept, 1980. Ploughed: 29 Dec. Spring-tine cultivated twice: 19 Feb, 1981, 20 Feb. Seed sown: 20 Feb. Weedkiller applied: 8 Apr. Fungicide applied: 18 June. Combine harvested: 9 Sept. Previous crops: W. wheat 1979, s. barley 1980.

NOTE: Leaf notching by Sitona was counted in June, and aphids were counted seven times during June and July.

81/R/BE/19

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
NONE	3.45
EL V2 D1	3.79
EL V2 D2	3.61
EL V2 D4	3.89
EL V2 P4	3.41
EL - U	3.00
EL V1 U	3.40 3.20
EL V2 U	
ST - D4	3.89
ST - P4	3.67
MFAN	3.53

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	TREATMNT
SED	0.287

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV% BLOCK.WP 27 0.406 11.5

GRAIN MEAN DM% 87.2

SUB PLOT AREA HARVESTED 0.00195