

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1981

[Full Table of Content](#)



## 81/W/RN/6 Arable Reference Plots - W. Barley, W. Oats

**Rothamsted Research**

Rothamsted Research (1982) *81/W/RN/6 Arable Reference Plots - W. Barley, W. Oats ; Yields Of The Field Experiments 1981*, pp 70 - 75 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/W/RN/6

ARABLE REFERENCE PLOTS

Object: To study the long term effects of FYM and N, P and K fertilisers on the yield and mineral content of crops - Woburn Stackyard C.

Sponsor: F.V. Widdowson.

The 22nd year, w. barley, w. oats.

For previous years see 60/B/3(t), 61-65/B/2, 66/B/2(t), 67/B/2(t), 68/B/3(t), 69/W/RN/6, 70/W/RN/6(t) and 71-80/W/RN/6.

Design: Blocks of 12 plots for each crop. Barley and oats one third replicate of 3 x 12 x 3 in 3 blocks of 12 plots.

Whole plot dimensions: 2.74 x 2.13.

Treatments: All combinations of:-

Blocks

1. CROP                   Crops in 1981 and previous crops:

W. oats after w. barley in 1980:

OW(OW)	After w. oats in 1979 (arable since 1960)
OW(BS)	After s. barley in 1979 (arable since 1960)
OW(BS G)	After s. barley in 1979 (grass 1960-73, arable thereafter)

W. barley after w. oats in 1980:

BW(P)	After potatoes in 1979
BW(SB)	After sugar beet in 1979
BW(L)	After one-year ley in 1979

Plots

2. MANURE(79)           Fertilisers and farmyard manure (until 1979, none in 1980 & 1981):

O  
N1  
P  
N1P  
K  
N1K  
PK  
N1PK  
N2PK  
D  
N1PKD  
N2PKD

N1,2 (kg N): 31.5, 63 (ley): 63, 126 (s. barley and w. oats): 126, 252 (sugar beet and potatoes): 188, 376 (permanent grass) as ammonium nitrate.

P: P205 at 63 kg as triple superphosphate.

81/W/RN/6

K: K20 at 252 kg as potassium bicarbonate.

D: Farmyard manure at 25 tonnes (permanent grass): 50 tonnes (sugar beet and potatoes): none to other crops.

and some of the combinations of 1 & 2 with:

3. N 81              Nitrogen fertiliser (kg N) (cumulative to 1980):

90	25 kg N to seedbed, 30 kg N in March, 35 kg N in April
120	25 kg N to seedbed, 40 kg N in March, 55 kg N in April
150	25 kg N to seedbed, 50 kg N in March, 75 kg N in April

NOTE: The block in permanent grass was discontinued in 1981.

Standard applications:

Both crops: Weedkillers: Ioxynil at 0.32 kg and mecoprop at 0.94 kg in 280 l, ioxynil at 0.42 kg and mecoprop at 1.3 kg in 280 l, on both occasions with tridemorph. Fungicides: Tridemorph at 0.53 kg, benomyl at 0.28 kg in 280 l.

W. oats: Insecticides: Phorate at 2 kg as granules, pirimicarb at 0.14 kg in 280 l. Fungicides: Carbendazim, tridemorph and maneb (as 'Cosmic' at 3.9 kg) with the pirimicarb.

W. barley: Fungicides: Benodanil at 1.1 kg with tridemorph at 0.53 kg in 280 l.

Seed: W. oats: Pennal, sown at 210 kg.

W. barley: Igri, dressed with ethirimol, sown at 190 kg.

Cultivations, etc.:-

W. oats and w. barley: Rotary cultivated: 15 Sept, 1980. Weedkillers and tridemorph applied: 25 Nov, 26 Mar, 1981. Second N applied: 18 Mar. Benomyl applied: 26 Mar. Third N applied: 24 Apr.

W. barley only: First N applied, seed sown: 15 Sept, 1980. Benodanil with tridemorph applied: 13 May, 1981. Harvested: 21 July.

W. oats: First N applied, phorate applied, raked in, seed sown: 2 Oct, 1980. 'Cosmic' and pirimicarb applied: 1 June, 1981. Harvested: 11 Aug.

NOTE: N, P and K contents of grain were measured.

81/W/RN/6

WINTER BARLEY

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CROP MANURE(79)	BW(P)	BW(SB)	BW(L)	MEAN
0	3.76	4.45	3.37	3.86
N1	3.24	3.25	2.20	2.90
P	2.85	4.33	1.33	2.84
N1P	1.57	3.65	1.51	2.25
K	6.86	6.13	5.98	6.32
N1K	7.16	6.00	5.40	6.19
PK	6.91	6.20	6.43	6.51
N1PK	6.75	6.12	6.35	6.41
N2PK	5.54	7.25	5.60	6.13
D	6.11	7.07	6.12	6.43
N1PKD	5.92	6.69	4.98	5.86
N2PKD	6.23	5.99	5.17	5.80
MEAN	5.24	5.59	4.54	5.12
N 81 MANURE(79)	90	120	150	MEAN
0	4.45	3.76	3.37	3.86
N1	2.20	3.25	3.24	2.90
P	4.33	2.85	1.33	2.84
N1P	1.51	3.65	1.57	2.25
K	6.13	6.86	5.98	6.32
N1K	5.40	6.00	7.16	6.19
PK	6.20	6.91	6.43	6.51
N1PK	6.35	6.12	6.75	6.41
N2PK	5.54	5.60	7.25	6.13
D	6.11	6.12	7.07	6.43
N1PKD	5.92	4.98	6.69	5.86
N2PKD	6.23	5.17	5.99	5.80
MEAN	5.03	5.11	5.24	5.12

GRAIN MEAN DM% 83.4

81/W/RN/6

WINTER BARLEY

STRAW TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CROP MANURE(79)	BW(P)	BW(SB)	BW(L)	MEAN
0	5.04	5.01	5.33	5.13
N1	4.40	4.48	3.72	4.20
P	6.19	5.40	3.92	5.17
N1P	5.20	4.92	2.68	4.27
K	7.69	6.21	8.13	7.35
N1K	7.81	7.39	6.66	7.29
PK	8.20	6.95	8.87	8.01
N1PK	9.08	7.62	7.63	8.11
N2PK	6.53	8.46	7.90	7.63
D	7.23	8.57	8.26	8.02
N1PKD	7.49	9.44	7.28	8.07
N2PKD	7.77	9.38	6.88	8.01
MEAN	6.89	6.99	6.44	6.77
N 81 MANURE(79)	90	120	150	MEAN
0	5.01	5.04	5.33	5.13
N1	3.72	4.48	4.40	4.20
P	5.40	6.19	3.92	5.17
N1P	2.68	4.92	5.20	4.27
K	6.21	7.69	8.13	7.35
N1K	6.66	7.39	7.81	7.29
PK	6.95	8.20	8.87	8.01
N1PK	7.63	7.62	9.08	8.11
N2PK	6.53	7.90	8.46	7.63
D	7.23	8.26	8.57	8.02
N1PKD	7.49	7.28	9.44	8.07
N2PKD	7.77	6.88	9.38	8.01
MEAN	6.11	6.82	7.38	6.77

STRAW MEAN DM% 72.1

PLOT AREA HARVESTED 0.00050

81/W/RN/6

WINTER OATS

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CROP MANURE(79)	OW(OW)	OW(BS)	OW(BS G)	MEAN
0	4.57	2.86	3.90	3.78
N1	3.06	3.24	3.94	3.41
P	4.42	2.56	2.69	3.23
N1P	2.29	3.36	4.07	3.24
K	4.68	4.31	5.85	4.95
N1K	4.25	4.58	4.73	4.52
PK	5.29	5.19	5.15	5.21
N1PK	3.53	4.71	4.19	4.14
N2PK	5.70	3.95	5.56	5.07
D	5.59	4.83	5.72	5.38
N1PKD	5.46	3.64	5.35	4.82
N2PKD	5.44	4.41	4.43	4.76
MEAN	4.52	3.97	4.63	4.38
N 81 MANURE(79)	90	120	150	MEAN
0	2.86	4.57	3.90	3.78
N1	3.94	3.24	3.06	3.41
P	2.56	4.42	2.69	3.23
N1P	4.07	3.36	2.29	3.24
K	4.31	4.68	5.85	4.95
N1K	4.73	4.58	4.25	4.52
PK	5.19	5.29	5.15	5.21
N1PK	4.19	4.71	3.53	4.14
N2PK	5.70	5.56	3.95	5.07
D	5.59	5.72	4.83	5.38
N1PKD	5.46	5.35	3.64	4.82
N2PKD	5.44	4.43	4.41	4.76
MEAN	4.50	4.66	3.96	4.38

GRAIN MEAN DM% 83.6

81/W/RN/6

WINTER OATS

STRAW TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CROP MANURE(79)	OW(OW)	OW(BS)	OW(BS G)	MEAN
0	5.94	2.96	5.91	4.94
N1	5.37	3.23	4.33	4.31
P	6.63	3.28	4.82	4.91
N1P	5.24	4.55	4.77	4.86
K	8.65	5.09	9.50	7.75
N1K	7.65	6.60	5.92	6.72
PK	8.57	7.43	9.66	8.56
N1PK	9.06	7.27	6.83	7.72
N2PK	8.97	6.70	7.99	7.89
D	8.30	10.51	11.38	10.06
N1PKD	8.68	8.49	11.37	9.52
N2PKD	8.88	9.33	10.17	9.46
MEAN	7.66	6.29	7.72	7.22
N 81 MANURE(79)	90	120	150	MEAN
0	2.96	5.94	5.91	4.94
N1	4.33	3.23	5.37	4.31
P	3.28	6.63	4.82	4.91
N1P	4.77	4.55	5.24	4.86
K	5.09	8.65	9.50	7.75
N1K	5.92	6.60	7.65	6.72
PK	7.43	8.57	9.66	8.56
N1PK	6.83	7.27	9.06	7.72
N2PK	8.97	7.99	6.70	7.89
D	8.30	11.38	10.51	10.06
N1PKD	8.68	11.37	8.49	9.52
N2PKD	8.88	10.17	9.33	9.46
MEAN	6.29	7.70	7.69	7.22

STRAW MEAN DM% 68.5

PLOT AREA HARVESTED 0.00050