

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1981

[Full Table of Content](#)



81/R/RN/7 Residual Phosphate - Ley

Rothamsted Research

Rothamsted Research (1982) *81/R/RN/7 Residual Phosphate - Ley* ; Yields Of The Field Experiments 1981, pp 76 - 79 - DOI: <https://doi.org/10.23637/ERADOC-1-35>

81/R/RN/7

RESIDUAL PHOSPHATE

Object: Originally to study the fresh and residual effects of phosphate fertiliser on the yields of three arable crops grown in rotation. From 1974 the effects on ley and on yield and pathogens of continuous w. wheat were studied. Since 1980 only ley has been included - Great Field IV and Sawyers I.

Sponsors: G.E.G. Mattingly.

The 22nd year, ley.

For previous years see 'Details' 1967 and 1973 and 74-80/R/RN/7.

Design: Great Field IV: 3 series each of 1 randomised block of 12 plots.
Sawyers I: 3 series each of 2 randomised blocks of 12 plots.

Whole plot dimensions:

Great Field IV: 4.27 x 18.3
Sawyers I: 4.27 x 20.1

Treatments:

P205 Rates and frequency of applying phosphate:-

NONE 0

Annual dressings, kg P205:

29 ANN	29
57 ANN	57
115 ANN	115
172 ANN	172

Triennial dressings, kg P205 (last applied 1978):

86 TRI	86
172 TRI	172

Six-yearly dressings, kg P205 (last applied 1973):

344 SIX	344
688 SIX	688
1032 SIX	1032

Single dressing, kg P205 (applied autumn 1959):

376 G(1)	376 as Gafsa rock phosphate
376 S(1)	376 as granular superphosphate

NOTES: (1) From 1974 the original rotation of potatoes, s. barley, swedes on both fields was changed. Blocks after barley were sown to w.wheat on Sawyers I, to ley on Great Field IV. In 1978 & 1979 one series was sown to ley each year on Sawyers I. In 1980 & 1981 the remaining series on Sawyers I was followed.

81/R/RN/7

- (2) Since 1960 all phosphate has been applied as superphosphate.
 (3) The six-yearly dressings were applied half in autumn before ploughing, half in spring.

Standard application:

Leys: Manures: K₂O at 250 kg as muriate of potash.

Cultivations, etc.:-

All Leys: K applied: 24 Nov, 1980.

Leys (Gt. Field IV only): Test P applied: 13 Feb. Cut: 1 June, 5 Aug, 4 Nov.

Leys (Sawyers I only): Test P applied: 25 Feb, 1981. Cut: 2 June, 7 Aug, 4 Nov.

Fallow: Ploughed: 8 Dec, 1980. Rotary cultivated: 16 June, 1981.

Spring-tine cultivated: 1 July.

GT. FIELD IV

SERIES I LEY

DRY MATTER TONNES/HECTARE

	CUT 1 (1/6/81)	CUT 2 (5/8/81)	CUT 3 (4/11/81)	TOTAL OF 3 CUTS
P205				
NONE	2.42	2.27	0.80	5.49
29 ANN	3.83	2.12	1.05	7.00
57 ANN	4.30	2.17	1.09	7.56
115 ANN	4.48	1.94	1.41	7.83
172 ANN	5.52	2.34	1.40	9.26
86 TRI	4.10	2.29	1.06	7.45
172 TRI	4.20	2.41	1.34	7.95
344 SIX	4.19	2.41	1.19	7.80
688 SIX	4.58	2.08	1.36	8.02
1032 SIX	4.67	2.59	1.48	8.74
376 G(1)	2.80	2.49	0.96	6.24
376 S(1)	2.89	2.51	0.93	6.34
MEAN	4.00	2.30	1.17	7.47
MEAN DM%	17.2	21.9	23.7	20.9

81/R/RN/7 GREAT FIELD IV

SERIES II LEY

DRY MATTER TONNES/HECTARE

	CUT 1 (1/6/81)	CUT 2 (5/8/81)	CUT 3 (4/11/81)	TOTAL OF 3 CUTS
P205				
NONE	1.47	2.65	0.76	4.88
29 ANN	4.15	2.59	1.34	8.08
57 ANN	4.77	2.62	1.15	8.54
115 ANN	4.65	2.63	1.23	8.50
172 ANN	5.10	3.20	1.33	9.63
86 TRI	2.97	2.79	1.04	6.80
172 TRI	4.76	2.89	1.12	8.78
344 SIX	3.78	2.78	1.09	7.65
688 SIX	5.01	2.98	1.38	9.37
1032 SIX	4.51	3.06	1.36	8.93
376 G(1)	2.06	2.83	0.87	5.77
376 S(1)	2.15	2.71	0.79	5.65
MEAN	3.78	2.81	1.12	7.71
MEAN DM%	8.6	23.7	21.8	21.4

SERIES III LEY

DRY MATTER TONNES/HECTARE

	CUT 1 (1/6/81)	CUT 2 (5/8/81)	CUT 3 (4/11/81)	TOTAL OF 3 CUTS
P205				
NONE	2.31	2.61	0.44	5.35
29 ANN	5.68	2.83	0.82	9.33
57 ANN	5.40	2.43	0.97	8.80
115 ANN	5.64	2.95	1.09	9.68
172 ANN	5.48	2.49	1.21	9.18
86 TRI	4.49	2.29	0.77	7.55
172 TRI	4.72	2.89	0.94	8.55
344 SIX	4.75	2.73	0.71	8.20
688 SIX	4.33	2.49	0.84	7.67
1032 SIX	5.10	2.41	0.95	8.47
376 G(1)	3.12	2.35	0.78	6.25
376 S(1)	1.95	2.70	0.52	5.17
MEAN	4.41	2.60	0.84	7.85
MEAN DM%	9.8	24.3	21.4	21.8

PLOT AREA HARVESTED 0.00186

81/R/RN/7 SAWYERS I

SERIES II LEY

DRY MATTER TONNES/HECTARE

	CUT 1 (2/6/81)	CUT 2 (7/8/81)	CUT 3 (4/11/81)	TOTAL OF 3 CUTS
P205				
NONE	3.68	1.25	0.66	5.58
29 ANN	4.51	1.48	0.68	6.68
57 ANN	4.67	1.39	0.69	6.76
115 ANN	5.41	1.18	0.63	7.22
172 ANN	5.95	1.05	0.54	7.54
86 TRI	4.14	1.31	0.58	6.03
172 TRI	4.47	1.45	0.68	6.61
344 SIX	4.35	1.49	0.64	6.48
688 SIX	4.39	1.52	0.70	6.61
1032 SIX	4.98	1.53	0.63	7.14
376 G(1)	3.65	1.35	0.43	5.43
376 S(1)	3.58	1.48	0.45	5.52
MEAN	4.48	1.37	0.61	6.47
SED*	0.288	0.123	0.102	0.410
CV%	6.4	8.9	16.7	6.3
MEAN DM%	18.4	18.4	22.2	19.7

* NOTE STRATUM STANDARD ERROR (11 df) is also equal to this figure

SERIES III LEY

DRY MATTER TONNES/HECTARE

	CUT 1 (2/6/81)	CUT 2 (7/8/81)	CUT 3 (4/11/81)	TOTAL OF 3 CUTS
P205				
NONE	3.29	1.21	0.59	5.08
29 ANN	5.03	1.04	0.31	6.38
57 ANN	5.37	1.18	0.34	6.89
115 ANN	5.40	0.90	0.27	6.57
172 ANN	5.53	0.97	0.15	6.66
86 TRI	4.74	0.97	0.28	5.99
172 TRI	4.85	0.97	0.36	6.19
344 SIX	4.68	1.01	0.44	6.12
688 SIX	5.21	1.07	0.37	6.65
1032 SIX	5.09	1.08	0.36	6.54
376 G(1)	3.72	1.12	0.44	5.28
376 S(1)	3.83	1.18	0.38	5.39
MEAN	4.73	1.06	0.36	6.14
SED*	0.170	0.057	0.073	0.125
CV%	3.6	5.4	20.4	2.0
MEAN DM%	19.5	20.3	23.0	20.9

PLOT AREA HARVESTED 0.00204