

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/W/CS/66 Dazomet and Nitrogen - Forage Maize

Rothamsted Research

Rothamsted Research (1983) *82/W/CS/66 Dazomet and Nitrogen - Forage Maize* ; Yields Of The Field Experiments 1982, pp 111 - 112 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/W/CS/66

DAZOMET AND NITROGEN

Object: To study the cumulative effects of dazomet and nitrogen on pathogens and yield of maize grown continuously - Woburn Butt Furlong.

Sponsors: A.J. Barnard, D. Hornby.

The 12th year, forage maize.

For previous years see 71/W/CS/66(t), 72/W/CS/66(t) and 73-81/W/CS/66.

Design: 2 blocks of 4 plots split into 4.

Whole plot dimensions: 2.13 x 16.5.

Treatments: All combinations of:-

Whole plots

1. DAZOMET(79) Dazomet (kg per annum) cumulative 1971-79, residual 1980-81:

0
450

2. DAZOMET(82) Dazomet (kg) in 1982 only:

0
450

Sub plots

3. N+FUNG Nitrogen fertilizer as 'Nitro-Chalk' and fungicide:

NONE	None
N78+N120	78 kg N on 5 Feb, 120 kg N to seedbed on 5 May
N120	120 kg N to seedbed on 5 May
N120+FOS	120 kg N to seedbed + 28 kg fosetyl-A1 to seedbed

NOTE: Sub plot treatments were superimposed on previous cumulative N treatments 1971-81.

Basal applications: Manures: (0:18:36) at 490 kg. Weedkiller: Atrazine at 1.1 kg in 280 l.

Seed: Fronica, sown at 103,000 seed per hectare.

Cultivations, etc.: - Ploughed: 16 Nov, 1981. Spring-tine cultivated: 24 Nov. Dazomet applied, rotary cultivated twice: 26 Nov. Early N applied: 5 Feb, 1982. Heavy spring-tine cultivated: 21 Apr. PK applied: 29 Apr. Seedbed N applied, fungicide treatment applied, rotary cultivated with crumbler attached, weedkiller applied, seed sown: 5 May. Missing portions of row resown by hand: 28 May. Hand harvested: 13 Oct.

NOTES (1): Soil samples were taken for estimates of total biomass.
(2): Plant samples were taken for assessments of bacteria and fungi on the roots.

82/W/CS/66

(3): Counts were made of common smut (*Ustilago maydis*) and stalk rots (*Fusarium* spp.)

FORAGE DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

DAZOMET(82)	0	450	MEAN		
DAZOMET(79)					
0	5.97	6.64	6.31		
450	6.64	6.85	6.74		
MEAN	6.30	6.75	6.53		
N+FUNG	NONE	N78+N120	N120	N120+FOS	MEAN
DAZOMET(79)					
0	3.13	7.81	7.11	7.16	6.31
450	3.29	8.06	8.13	7.49	6.74
MEAN	3.21	7.94	7.62	7.33	6.53
N+FUNG	NONE	N78+N120	N120	N120+FOS	MEAN
DAZOMET(82)					
0	2.92	8.24	7.09	6.97	6.30
450	3.50	7.64	8.16	7.69	6.75
MEAN	3.21	7.94	7.62	7.33	6.53
DAZOMET(79)	N+FUNG	NONE	N78+N120	N120	N120+FOS
0	0	3.08	7.60	6.62	6.58
	450	3.19	8.03	7.61	7.75
450	0	2.77	8.87	7.55	7.36
	450	3.81	7.25	8.72	7.63

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	N+FUNG	DAZOMET(79)* N+FUNG	DAZOMET(82)* N+FUNG	DAZOMET(79)* DAZOMET(82) N+FUNG
-------	--------	------------------------	------------------------	---------------------------------------

SED	0.315	0.445	0.445	0.629
-----	-------	-------	-------	-------

* WITHIN SAME LEVEL OF DAZOMET(79) OR DAZOMET(82) OR BOTH

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
---------	----	----	-----

BLOCK.WP.SP	12	0.629	9.6
-------------	----	-------	-----

GRAIN MEAN DM% 29.7

SUB PLOT AREA HARVESTED 0.00078