Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1982



Full Table of Content

82/R/CS/130 Effects of Earthworm Inoculation - Ley

Rothamsted Research

Rothamsted Research (1983) 82/R/CS/130 Effects of Earthworm Inoculation - Ley; Yields Of The Field Experiments 1982, pp 116 - 118 - DOI: https://doi.org/10.23637/ERADOC-1-33

82/R/CS/130

EFFECTS OF EARTHWORM INOCULATION

Object: To study the effects of different species of earthworms on yield and other characteristics of grass - Fosters O and E.

Sponsor: J.R. Lofty.

The ninth year, ley.

For previous years see 74-81/R/CS/130.

Design: 3 randomised blocks of 4 plots.

Whole plot dimensions: 8.53 x 9.14.

Treatments: Inoculation with earthworm species in 1974, 1975 and 1979:

WORMSPEC

NONE	None
ALLOLOB0	Allolobophora longa at 15,000 per hectare in 1974; 5,000 in 1975; 96,000 in 1979
LUMBRICU	Lumbricus terrestris at 5,000 per hectare in 1974 and 1975; 96,000 in 1979
SIX SPEC	Six species - A. caliginosa, A. chlorotica, A. longa, A. rosea, L. rubellus, L. terrestris at a total of 35,000

- NOTES: (1) The experiment was ploughed in error in July 1976 and resown in autumn 1976.
 - (2) Earthworms for the 1979 crop were applied in December, 1978 to one block only. Applications to other blocks have been postponed.

per hectare in 1974; 12,000 in 1975; none in 1979

- Basal applications: Manures: (0:14:28) at 500 kg. (25:0:16) at 440 kg in spring, (25:0:16) at 220 kg after the first two cuts.
- Seed: Grass and clover mixture sown August 1973, ploughed in error in June 1976 and resown September 1976.
- Cultivations, etc.:- PK applied: 13 Nov, 1981. NK applied: 22 Mar, 1982, 7 June, 16 July. Cut: 1 June, 14 July, 10 Nov.

82/R/CS/130

1ST CUT (1/6/82) DRY MATTER TONNES/HECTARE

**** TABLES OF MEANS *****

WORMSPEC

NONE ALLOLOBO LUMBRICU SIX SPEC

MEAN

3.71

4.05 3.63 3.66

3.76

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE

WORMSPEC

SED

0.407

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM

DF

SE

CV%

BLOCK. WP

6

0.498

13.2

1ST CUT MEAN DM% 23.7

2ND CUT (14/7/82)DRY MATTER TONNES/HECTARE

**** TABLES OF MEANS ****

WORMSPEC

1.81 1.98

NONE ALLOLOBO LUMBRICU SIX SPEC 1.91

MEAN 1.87

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE

WORMSPEC

SED

0.057

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM

DF

SE

CV%

BLOCK. WP

6

0.070

3.7

2ND CUT MEAN DM% 20.5

82/R/CS/130

3RD CUT (10/11/82)DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

WORMSPEC NONE ALLOLOBO LUMBRICU SIX SPEC MEAN 1.01 0.78 1.05 0.89 0.93

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE WORMSPEC
SED 0.170

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CY%

BLOCK.WP 6 0.208 22.3

3RD CUT MEAN DM% 36.0

TOTAL OF 3 CUTS DRY MATTER TONNES/HECTARE

**** TABLES OF MEANS *****

WORMSPEC NONE ALLOLOBO LUMBRICU SIX SPEC MEAN 6.53 6.80 6.59 6.34 6.56

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE WORMSPEC
SED 0.539

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV%
BLOCK.WP 6 0.660 10.1

TOTAL OF 3 CUTS MEAN DM% 26.7

PLOT AREA HARVESTED 0.00046