

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)

International Agricultural Sciences
Number
Year Published
Title
Author
Keywords
DOI
This document is licensed under a Creative Commons Attribution 4.0 International License and is available at <https://www.era-doc.org/>. For more information on the ERA-DOC project, please visit <https://www.era-doc.org/>.
157-159
1982

82/R/CS/264 Fungicide Rates - W. Barley, S. Barley

Rothamsted Research

Rothamsted Research (1983) *82/R/CS/264 Fungicide Rates - W. Barley, S. Barley* ; Yields Of The Field Experiments 1982, pp 157 - 159 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/CS/264

FUNGICIDE RATES

Object: To study the residual effects on barley of a range of rates of triadimefon applied to w. oilseed rape on land in rotation (pathogen free) and after previous oilseed rape (pathogen infected) - Long Hoos IV 1 (pathogen free) and Summerdells I (pathogen infected).

Sponsor: C.J. Rawlinson.

The second year, w. barley (Long Hoos IV 1), s. barley (Summerdells I).

For previous year see 81/R/CS/264.

Design: 3 randomised blocks of 6 plots.

Whole plot dimensions: 2.13 x 3.05.

Treatments:

FUNGRATE (81) Residues of triadimefon applied at different rates (kg) to soil surface before seedbed cultivation for w. oilseed rape 1981:

0.06
0.12
0.25
0.50
1.00
2.00

NOTE: In 1981 on Summerdells I w. oilseed rape failed during the winter and the site was sown to s. barley to assess the residual effects of treatments already applied.

Basal applications:

Long Hoos IV: Manures: 'Nitro-Chalk' at 500 kg. Weedkillers: Mecoprop, bromoxynil and ioxynil (as 'Brittox' at 2.8 l) in 450 l. Dicamba, mecoprop and MCPA (as 'Banlene Plus' at 4.2 l) in 340 l.

Summerdells I: Manures: (0:14:28) at 780 kg, 'Nitro-Chalk' at 470 kg. Weedkillers: Paraquat at 0.5 kg ion in 250 l. Dicamba, mecoprop and MCPA (as 'Poly-Farmon' at 5.0 l) in 250 l.

Seed: Long Hoos IV: Maris Otter, undressed, sown at 160 kg.
Summerdells I: Georgie, sown at 160 kg.

Cultivations, etc.:-

Long Hoos IV: Ploughed: 6 Aug, 1981. Power harrowed: 23 Sept. Spring-tine cultivated: 13 Oct. Seed sown: 16 Oct. Mecoprop, bromoxynil and ioxynil applied: 1 Feb, 1982. Dicamba, mecoprop and MCPA applied: 19 Apr. N applied: 20 Apr. Combine harvested: 26 July.

Summerdells I: Ploughed: 8 Oct, 1981. PK applied: 26 Nov. N and paraquat applied: 25 Mar, 1982. Spring-tine cultivated: 30 Mar. Rotary harrowed, seed sown: 3 Apr. Dicamba, mecoprop and MCPA applied: 17 May. Combine harvested: 17 Aug.

82/R/CS/264

NOTE: Mildew and other diseases were assessed five times at fortnightly intervals from May to July. Components of yield were measured at maturity.

LONG HOOS IV WINTER BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGRATE	0.06	0.12	0.25	0.50	1.00	2.00	MEAN
	2.81	2.59	2.17	2.36	1.67	2.90	2.42

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGRATE
-----	-----
SED	0.413

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.505	20.9

GRAIN MEAN DM% 69.2

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGRATE	0.06	0.12	0.25	0.50	1.00	2.00	MEAN
	5.57	5.42	5.65	6.20	6.26	5.54	5.77

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGRATE
-----	-----
SED	0.875

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	10	1.072	18.6

STRAW MEAN DM% 58.8

PLOT AREA HARVESTED 0.00054

82/R/CS/264

SUMMERDELLS I SPRING BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGRATE	0.06	0.12	0.25	0.50	1.00	2.00	MEAN
	4.68	4.35	4.58	4.66	4.22	4.88	4.56

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGRATE
-----	-----
SED	0.448

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.549	12.0

GRAIN MEAN DM% 82.0

PLOT AREA HARVESTED 0.00065