

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1982

[Full Table of Content](#)



## 82/R/CS/279 Nematicides and Stem Nematode

### Rothamsted Research

Rothamsted Research (1983) *82/R/CS/279 Nematicides and Stem Nematode ; Yields Of The Field Experiments 1982*, pp 174 - 180 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/CS/279

NEMATICIDES AND STEM NEMATODE

Object: To study, on sites free from or infested by stem nematode (*Ditylenchus dipsaci*), the effects of nematicides on lucerne - Long Hoos V 5 (healthy) and Long Hoos IV 2 (infested).

Sponsor: A.G. Whitehead.

The first year, lucerne.

Design: On each site: 3 randomised blocks of 14 plots.

Whole plot dimensions: 1.22 x 3.66.

Treatments (applied to HEALTHY AND INFESTED sites):

TREATMNT	Varieties, rates and methods of applying nematicides:
V 0	Vertus, untreated
V A1	Vertus, aldicarb at 1.5 kg in seed furrow
E 0	Europe, untreated
E A1	Europe, aldicarb at 1.5 kg in seed furrow (duplicated)
E A2	Europe, aldicarb at 3.0 kg in seed furrow (duplicated)
E A1 A1	Europe, aldicarb at 1.5 kg in seed furrow, repeated after each cut
E C1	Europe, carbofuran at 1.5 kg in seed furrow (duplicated)
E C2	Europe, carbofuran at 3.0 kg in seed furrow (duplicated)
E T1	Europe, thiabendazole at 1.5 kg over the rows at drilling
E T2	Europe, thiabendazole at 3.0 kg over the rows at drilling

Basal applications: Manures: (0:25:25) at 530 kg. Weedkiller: 2.4 - DB at 2.8 l in 280 l.

Seed: Both sites: Varieties sown at 11 kg.

Cultivations, etc.:-

HEALTHY site: PK applied: 26 Mar, 1982. Rotary harrowed twice, Europe sown, aldicarb and carbofuran applied to this variety: 29 Mar. Vertus sown and aldicarb applied to this variety, thiabendazole applied to Europe: 30 Mar. Weedkiller applied: 26 May. Cut: 13 July. Aldicarb applied: 16 July. Cut, aldicarb applied: 26 Aug. Previous crops: Potatoes 1980, s.wheat 1981.

INFESTED site: Ploughed: 24 Sept, 1981. PK applied: 26 Mar, 1982. Rotary harrowed, seed sown, aldicarb, carbofuran and thiabendazole applied to Europe and aldicarb to Vertus: 31 Mar. Weedkiller applied: 26 May. Cut north block: 13 July. Aldicarb applied to this block: 16 July. Cut remaining two blocks: 19 July. Aldicarb applied to these two blocks: 22 July. Cut: 25 Aug. Aldicarb applied: 26 Aug. Previous crops: Fallow 1980, lucerne 1981.

82/R/CS/279 LONG HOOS V 5 (HEALTHY SITE)  
 1ST CUT (13/7/82) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT	
V 0	2.04
V A1	1.38
E 0	2.91
E A1	2.72
E A2	2.41
E A1 A1	2.68
E C1	3.55
E C2	3.25
E T1	3.11
E T2	2.84
MEAN	2.77

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT
-----	-----
SED	0.202 MIN REP
	0.175 MAX-MIN
	0.143 MAX REP

TREATMNT  
 MAX REP E A1 V E A2 V E C1 OR E C2  
 MAX-MIN E A1 V E A2 V E C1 OR E C2 V ANY OF REMAINDER  
 MIN REP ANY OF REMAINDER

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	30	0.247	8.9
1ST CUT MEAN DM%	17.8		

82/R/CS/279 LONG HOOS V 5 (HEALTHY SITE)

2ND CUT (26/8/82) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT	
V 0	2.75
V A1	2.20
E 0	3.15
E A1	2.90
E A2	2.89
E A1 A1	3.16
E C1	3.61
E C2	3.38
E T1	3.51
E T2	2.96
MEAN	3.09

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT
-----	-----
SED	0.208 MIN REP
	0.181 MAX-MIN
	0.147 MAX REP

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	30	0.255	8.3
2ND CUT MEAN DM%	21.9		

82/R/CS/279 LONG HOOS V 5 (HEALTHY SITE)

TOTAL OF 2 CUTS DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT	
V 0	4.79
V A1	3.58
E 0	6.05
E A1	5.62
E A2	5.30
E A1 A1	5.84
E C1	7.16
E C2	6.63
E T1	6.62
E T2	5.79
MEAN	5.87

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT
-----	-----
SED	0.360 MIN REP
	0.312 MAX-MIN
	0.255 MAX REP

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	30	0.441	7.5
TOTAL OF 2 CUTS MEAN DM%	19.9		
PLOT AREA HARVESTED	0.00045		

82/R/CS/279 LONG HOOS IV 2 (INFESTED SITE)

1ST CUT (13-19/7/82) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT	
V 0	2.42
V A1	1.89
E 0	3.49
E A1	3.66
E A2	3.27
E A1 A1	3.73
E C1	4.12
E C2	4.03
E T1	3.12
E T2	3.66
MEAN	3.46

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT
-----	-----
SED	0.224 MIN REP
	0.194 MAX-MIN
	0.159 MAX REP

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	30	0.275	7.9
1ST CUT MEAN DM%	21.6		



82/R/CS/279 LONG HOOS IV 2 (INFESTED SITE)  
2ND CUT (25/8/82) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT	
V 0	2.16
V A1	1.99
E 0	2.62
E A1	2.62
E A2	2.57
E A1 A1	2.70
E C1	2.92
E C2	2.80
E T1	2.53
E T2	2.61
MEAN	2.60

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT
-----	-----
SED	0.117 MIN REP
	0.101 MAX-MIN
	0.083 MAX REP

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	30	0.143	5.5
2ND CUT MEAN DM%		24.2	

82/R/CS/279 LONG HOOS IV 2 (INFESTED SITE)

TOTAL OF 2 CUTS DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT	
V 0	4.58
V A1	3.88
E 0	6.11
E A1	6.27
E A2	5.83
E A1 A1	6.43
E C1	7.03
E C2	6.83
E T1	5.64
E T2	6.26
MEAN	6.06

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT
SED	0.273 MIN REP
	0.236 MAX-MIN
	0.193 MAX REP

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	30	0.334	5.5
TOTAL OF 2 CUTS MEAN DM%	22.9		
PLOT AREA HARVESTED	0.00045		