Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1982



Full Table of Content

82/R/WW/5 Nuarimol and Take-all - Winter Wheat

Rothamsted Research

Rothamsted Research (1983) 82/R/WW/5 Nuarimol and Take-all - Winter Wheat; Yields Of The Field Experiments 1982, pp 236 - 237 - DOI: https://doi.org/10.23637/ERADOC-1-33

82/R/WW/5

WINTER WHEAT

NUARIMOL AND TAKE-ALL

Object: To study the effects of nuarimol on the incidence of take-all and on the yield of w. wheat - New Zealand.

Sponsor: G.L. Bateman.

Design: 5 randomised blocks of 5 plots.

Whole plot dimensions: 2.13 x 12.2.

Treatments:

| FUN CULT | Fungicide and methods of incorporation: |
|----------|---|
| NONE | None |
| NU EC PL | Nuarimol e.c. applied to stubble 1981 and ploughed in |
| NU EC LH | Nuarimol e.c. lightly rotary harrowed into seedbed |
| NU EC DC | Nuarimol e.c. deeply rotary cultivated into seedbed |
| NU WP DC | Nuarimol w.p. deeply rotary cultivated into seedbed |

NOTE: Nuarimol was applied at 2.2 kg.

Basal applications: Manures: (0:14:28) at 320 kg. 'Nitro-Chalk' at 560 kg. Weedkillers: Chlortoluron at 5.6 l in 250 l. Glyphosate at 1.4 kg in 250 l. Fungicides: Carbendazim, maneb and tridemorph (as 'Cosmic' at 4.0 kg) with captafol at 1.2 kg applied with the insecticide in 250 l. Insecticide: Pirimicarb at 0.14 kg. Growth regulator: Chlormequat at 1.7 kg in 250 l.

Seed: Avalon, untreated, sown at 200 kg.

Cultivations, etc.:- Ploughed: 1 Oct, 1981. PK applied: 26 Oct. Rotary harrowed, seed sown: 28 Oct. Chlortoluron applied: 29 Oct. N applied: 22 Apr, 1982. Growth regulator applied: 27 Apr. Fungicides and insecticide applied: 15 June. Glyphosate applied: 10 Aug. Combine harvested: 21 Aug. Previous crops: W. wheat 1980 and 1981.

NOTE: Take-all and foot rots were assessed in early May, early June and mid-July.

82/R/WW/5

GRAIN TONNES/HECTARE

**** TABLES OF MEANS *****

FUN CULT NONE NU EC PL NU EC LH NU EC DC NU WP DC MEAN 9.23 9.12 9.11 9.15 9.38 9.20

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE FUN CULT
SED 0.207

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV%

BLOCK.WP 16 0.328 3.6

GRAIN MEAN DM% 84.6

PLOT AREA HARVESTED 0.00176