

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/WW/6 Fungicides and Microflora - Winter Wheat

Rothamsted Research

Rothamsted Research (1983) *82/R/WW/6 Fungicides and Microflora - Winter Wheat* ; Yields Of The Field Experiments 1982, pp 238 - 240 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/WW/6

WINTER WHEAT

FUNGICIDES AND MICROFLORA

Object: To study the effects of a range of fungicides and times of application on the colonisation of fungi on flag leaves and developing ears and on the yield of w. wheat - Geescroft.

Sponsor: N. Magan.

Design: 2 randomised blocks of 24 plots.

Whole plot dimensions: 3.0 x 14.0.

Treatments: All combinations of:-

1. E FUNG Early-applied fungicides:

NONE	None
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg applied on 26 May

2. L FUNG A Late-applied fungicides:

BE+MN+MZ	Benomyl at 0.56 kg, plus maneb at 0.78 kg, and mancozeb at 0.78 kg
CAPTAFOL	Captafol at 1.4 kg
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg
IMAZALIL	Imazalil at 0.39 kg
PROCHLOR	Prochloraz at 0.40 l

3. LFNGDATE Dates of applying late fungicide:

M	10 June, 1982
L	16 June

plus two extra treatments not given L FUNG:

L FUNG 0

NONE	No early-applied fungicide (duplicated)
CARB+MAN	Carbendazim at 0.25 kg plus maneb at 1.6 kg applied on 26 May (duplicated)

NOTE: Treatment sprays were applied in 340 l.

Basal applications: Manures: 'Nitro-Chalk' applied at 480 kg. Weedkillers: Mecoprop (as 'Methoxone' at 5.0 l) with isoproturon at 2.0 kg in 250 l.

Seed: Maris Huntsman, sown at 190 kg.

Cultivations, etc.: - Spring-tine cultivated: 8 Oct, 1981. Heavy spring-tine cultivated twice: 12 Oct. Spring-tine cultivated, rotary harrowed, seed sown: 13 Oct. Weedkillers applied: 15 Apr, 1982. N applied: 23 Apr. Combine harvested: 11 Aug. Previous crops: W. beans 1980, potatoes 1981.

82/R/WW/6

NOTE: Grain and leaf microflora, especially Alternaria and Cladosporium, were assessed fortnightly from June to August.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

L FUNG A	BE+MN+MZ	CAPTAFOL	CARB+MAN	IMAZALIL	PROCHLOR	MEAN
E FUNG						
NONE	8.14	8.37	7.98	7.63	7.98	8.02
CARB+MAN	8.64	8.17	8.45	8.29	8.39	8.39
MEAN	8.39	8.27	8.22	7.96	8.19	8.20
LFNGDATE	M	L	MEAN			
E FUNG						
NONE	8.16	7.89	8.02			
CARB+MAN	8.38	8.39	8.39			
MEAN	8.27	8.14	8.20			
LFNGDATE	M	L	MEAN			
L FUNG A						
BE+MN+MZ	8.20	8.58	8.39			
CAPTAFOL	8.30	8.25	8.27			
CARB+MAN	8.38	8.05	8.22			
IMAZALIL	8.07	7.84	7.96			
PROCHLOR	8.40	7.97	8.19			
MEAN	8.27	8.14	8.20			
E FUNG	LFNGDATE	M	L			
NONE	L FUNG A					
	BE+MN+MZ	7.91	8.37			
	CAPTAFOL	8.50	8.25			
	CARB+MAN	8.32	7.64			
	IMAZALIL	7.65	7.60			
	PROCHLOR	8.39	7.58			
CARB+MAN	BE+MN+MZ	8.48	8.80			
	CAPTAFOL	8.09	8.25			
	CARB+MAN	8.44	8.46			
	IMAZALIL	8.49	8.08			
	PROCHLOR	8.42	8.36			
L FUNG O	NONE	CARB+MAN	MEAN			
	8.07	7.75	7.91			
GRAND MEAN	8.16					

82/R/WW/6

GRAIN TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	L FUNG O	E FUNG	L FUNG A	LFNGDATE
SED	0.364	0.163	0.258	0.163

TABLE	E FUNG L FUNG A	E FUNG LFNGDATE	L FUNG A LFNGDATE	E FUNG L FUNG A LFNGDATE
SED	0.364	0.230	0.364	0.515

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	25	0.515	6.3

GRAIN MEAN DM% 85.9

PLOT AREA HARVESTED 0.00204