Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1982



Full Table of Content

82/R/WW/14 - Winter Wheat

Rothamsted Research

Rothamsted Research (1983) 82/R/WW/14 - Winter Wheat; Yields Of The Field Experiments 1982, pp 243 - 244 - DOI: https://doi.org/10.23637/ERADOC-1-33

82/R/WW/14

WINTER WHEAT

Object: To study the effects of methods of applying benomyl on the incidence of eyespot and on the yield of w. wheat - Meadow.

Sponsors: G.R. Cayley, D.C. Griffiths, T. Fox.

Design: 4 randomised blocks of 3 plots.

Whole plot dimensions: 3.0×10.0 .

Treatments:

SPRAYERS

Sprayers used to apply benomyl:

NONE ELECSTAT HYDRAUL No spray applied Electrostatic sprayer

Standard hydraulic sprayer

NOTE: Benomyl was applied at 0.25 kg on 21 Apr, 1982, in 4.6 l for ELECTSTAT, in 490 l for HYDRAUL.

Basal applications: Manures: (0:14:28) at 320 kg. 'Nitro-Chalk' at 670 kg. Weedkillers: Mecoprop, bromoxynil and ioxynil (as 'Brittox' at 3.5 1) with isoproturon at 2.0 kg in 250 l.

Seed: Avalon, sown at 190 kg.

Cultivations, etc.:- Ploughed: 17 Oct, 1981. PK applied, rotary harrowed, seed sown: 26 Oct. Weedkillers applied: 14 Apr, 1982. N applied: 22 Apr. Combine harvested: 11 Aug. Previous crops: W. wheat 1980 and 1981.

NOTES: Eyespot (Pseudocercosporella herpotrichoides) and sharp eyespot (Rhizoctonia cerealis) were assessed and plant samples were analysed for chemical residues.

82/R/WW/14

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

SPRAYERS

NONE ELECSTAT HYDRAUL 5.85 6.04

6.02 6.17

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE

SPRAYERS

SED

0.157

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM

DF

SE

CV%

BLOCK. WP

6

0.222 3.7

GRAIN MEAN DM% 86.6

PLOT AREA HARVESTED 0.00204