

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1982

[Full Table of Content](#)



## 82/R/WW/14 - Winter Wheat

### Rothamsted Research

Rothamsted Research (1983) *82/R/WW/14 - Winter Wheat* ; Yields Of The Field Experiments 1982, pp 243 - 244 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/WW/14

WINTER WHEAT

Object: To study the effects of methods of applying benomyl on the incidence of eyespot and on the yield of w. wheat - Meadow.

Sponsors: G.R. Cayley, D.C. Griffiths, T. Fox.

Design: 4 randomised blocks of 3 plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments:

SPRAYERS	Sprayers used to apply benomyl:
NONE	No spray applied
ELECSTAT	Electrostatic sprayer
HYDRAUL	Standard hydraulic sprayer

NOTE: Benomyl was applied at 0.25 kg on 21 Apr, 1982, in 4.6 l for ELECTSTAT, in 490 l for HYDRAUL.

Basal applications: Manures: (0:14:28) at 320 kg. 'Nitro-Chalk' at 670 kg. Weedkillers: Mecoprop, bromoxynil and ioxynil (as 'Brittox' at 3.5 l) with isoproturon at 2.0 kg in 250 l.

Seed: Avalon, sown at 190 kg.

Cultivations, etc.: - Ploughed: 17 Oct, 1981. PK applied, rotary harrowed, seed sown: 26 Oct. Weedkillers applied: 14 Apr, 1982. N applied: 22 Apr. Combine harvested: 11 Aug. Previous crops: W. wheat 1980 and 1981.

NOTES: Eyespot (*Pseudocercospora herpotrichoides*) and sharp eyespot (*Rhizoctonia cerealis*) were assessed and plant samples were analysed for chemical residues.

82/R/WW/14

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

SPRAYERS	NONE	ELECSTAT	HYDRAUL	MEAN
	5.85	6.04	6.17	6.02

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	SPRAYERS
-----	-----
SED	0.157

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	6	0.222	3.7

GRAIN MEAN DM% 86.6

PLOT AREA HARVESTED 0.00204