

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/AG/6 Agdell - W. Beans, W. Wheat

Rothamsted Research

Rothamsted Research (1983) *82/R/AG/6 Agdell - W. Beans, W. Wheat* ; Yields Of The Field Experiments 1982, pp 26 - 30 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/AG/6

AGDELL

Object: To study, by crop yields and soil analyses, the residual values of phosphate and potash applied in the period 1848-1951 and further dressings since 1964.

The 13th year of revised scheme, w. beans, w. wheat.

For previous years see 'Details' 1967 and 1973, and 74-81/R/AG/6.

Treatments: All combinations of:-

Whole plots

1. OLDRESD Fertilizers and organic manures applied to roots every fourth year, in the period 1848-1948:

NONE	None
PKNAMG	P K Na Mg
NPKNAMGC	N P K Na Mg C

N: 48 kg N as sulphate of ammonia
P: 41 kg P as superphosphate
K: 224 kg K as sulphate of potash
Na: 16 kg Na as sulphate of soda
Mg: 11 kg Mg as sulphate of magnesia
C: Castor meal at 2240 kg supplying about 112 kg N

2. RN CROP Rotation 1848-1951 and crop in 1982:

F/WHEAT With fallow: Roots (turnips or swedes), s. barley, fallow, w. wheat 1848-1951. W. wheat (after w.beans 1981)

L/BEANS With legume: Roots, s. barley, legume (clover or beans), w. wheat 1848-1951. W. beans (after w. wheat 1981)

Half plots

3. 1964RESID Residues of 1964 treatments:

P
K

Quarter plots

4. PREVCROP Previous cropping 1958-69 on P-test half plots, 1958-70 on K-test half plots:

ARABLE	Arable or fallow
GRASS	Grass

82/R/AG/6

Sixteenth plots

5. P ₂ O ₅ 64	K ₂ O 64	Rates of 1964 treatments (kg):	
		P ₂ O ₅ to P-test half plots	K ₂ O to K-test half plots
0	0		
500	315		
1000	630		
2000	1260		

Thirty second plots

6.		On P-test half plots: To RN CROP F/WHEAT. Residues of P ₂ O ₅ applied 1970-72 (kg) and in 1979 and 1981 (kg):	
P ₂ O ₅ 721			
(0)0		None	
(375)300		375 total in 1970-72, 150 in 1979, 150 in 1981	
		To RN CROP L/BEANS. Residues of P ₂ O ₅ applied 1970-72 (kg) and fresh dressings in 1980 and 1982 (kg):	
P ₂ O ₅ 722			
(0)0		None	
(375)300		375 total in 1970-72, 150 in 1980, 150 in 1982	
		On K-test half plots: To RN CROP F/WHEAT. Residues of K ₂ O applied 1973-76 (kg) and in 1979 and 1981 (kg):	
K ₂ O 761			
(0)0		None	
(870)600		870 total in 1973-76, 300 in 1979, 300 in 1981	
		To RN CROP L/BEANS. Residues of K ₂ O applied 1973-76 (kg) and fresh dressings in 1980 and 1982 (kg):	
K ₂ O 762			
(0)0		None	
(870)600		870 total in 1973-76, 300 in 1980, 300 in 1982	

Standard applications:

Both crops: Weedkillers: Glyphosate at 1.4 kg in 250 l. Paraquat at 0.3 kg ion and diquat at 0.3 kg ion in 250 l.
W. beans: Weedkillers: Propyzamide at 0.85 kg in 250 l. Diquat at 0.42 kg ion (applied with 'Agral', a wetting agent) in 250 l.
Nematicide: Aldicarb at 10 kg.

82/R/AG/6

W. wheat: Manures: 'Nitro-Chalk' at 130 kg and 770 kg. Weedkillers: Chlortoluron at 5.6 l in 250 l. Mecoprop, bromoxynil and ioxynil (as 'Brittox' at 3.5 l) in 250 l. Glyphosate at 1.4 kg in 250 l. Fungicide: Propiconazole at 0.12 kg in 250 l and on a second occasion at 0.25 kg in 250 l with the insecticide. Insecticide: Pirimicarb at 0.14 kg.

Seed: W. beans: Throws MS, sown at 300 kg.
W. wheat: Avalon, sown at 200 kg.

Cultivations, etc.:-

Both crops: Glyphosate applied: 16 Sept, 1981. Paraquat and diquat applied: 7 Oct.

W. beans: Aldicarb applied, seed direct drilled: 29 Oct, 1981. Propyzamide applied: 30 Oct. Diquat applied: 25 Aug, 1982. Combine harvested: 2 Sept.

W. wheat: Seed direct drilled: 13 Oct, 1981. Chlortoluron applied: 16 Oct. N applied: 25 Feb, 1982 and 23 Apr. Mecoprop, bromoxynil and ioxynil applied: 19 Apr. Propiconazole applied: 26 May. Propiconazole with insecticide applied: 15 June. Glyphosate applied: 4 Aug. Combine harvested: 21 Aug.

82/R/AG/6

WHEAT P PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	OLDRES	NONE	PKNAMG		NPKNAMGC		
	P205 722	(0)0	(375)300	(0)0	(375)300	(0)0	(375)300
PREVCROP	P205 64						
ARABLE	0	6.51	7.73	7.06	6.26	3.88	3.80
	500	7.57	8.16	7.98	8.57	5.58	5.48
	1000	7.97	7.79	6.04	7.07	5.97	4.31
	2000	7.73	8.22	8.26	8.66	6.43	7.68
GRASS	0	5.27	7.44	4.65	7.37	7.06	7.73
	500	6.78	6.03	7.22	8.06	6.52	7.16
	1000	6.93	8.32	7.66	7.37	5.37	6.90
	2000	6.90	7.44	8.42	8.76	5.13	5.92

GRAIN MEAN DM% 83.6

WHEAT K PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	OLDRES	NONE	PKNAMG		NPKNAMGC		
	K20 762	(0)0	(870)600	(0)0	(870)600	(0)0	(870)600
PREVCROP	K20 64						
ARABLE	0	7.35	8.36	8.41	8.08	7.55	7.89
	315	7.28	7.95	8.68	8.75	8.16	8.20
	630	7.99	8.56	6.40	7.16	7.87	7.68
	1260	7.85	8.14	7.15	6.87	7.75	7.37
GRASS	0	4.73	6.81	7.73	8.43	6.97	7.98
	315	6.78	7.74	8.09	8.11	7.48	8.62
	630	7.70	8.23	7.98	8.34	6.84	7.77
	1260	7.50	8.34	8.24	8.54	6.68	7.33

GRAIN MEAN DM% 83.7

PLOT AREA HARVESTED (OLDRES NONE) 0.00121

PLOT AREA HARVESTED (REMAINDER) 0.00136

82/R/AG/6

BEANS P PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	OLDRESD	NONE	PKNAMG		NPKNAMGC		
	P205 721	(0)0	(375)300	(0)0	(375)300	(0)0	(375)300
PREVCROP	P205 64						
ARABLE	0	0.92	2.73	1.78	0.76	1.53	1.83
	500	1.98	2.93	2.37	1.95	0.47	0.54
	1000	2.48	2.99	0.45	0.99	1.61	1.53
	2000	1.40	3.57	1.60	2.57	0.85	1.00
GRASS	0	2.36	1.36	0.69	0.61	1.89	2.63
	500	1.87	0.85	0.11	0.53	1.88	1.29
	1000	1.61	0.85	0.46	0.61	2.10	1.29
	2000	2.62	1.11	0.76	0.39	2.82	2.94

GRAIN MEAN DM% 80.9

BEANS K PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	OLDRESD	NONE	PKNAMG		NPKNAMGC		
	K20 761	(0)0	(870)600	(0)0	(870)600	(0)0	(870)600
PREVCROP	K20 64						
ARABLE	0	2.01	3.07	2.50	2.52	1.22	1.37
	315	3.36	3.69	2.56	3.11	0.70	0.46
	630	2.76	2.90	2.29	2.17	1.91	1.59
	1260	3.32	2.82	2.67	3.34	1.08	0.77
GRASS	0	2.61	2.11	1.15	1.22	1.04	2.44
	315	2.84	2.36	1.12	2.22	1.05	1.86
	630	2.87	1.61	2.06	0.76	0.38	0.76
	1260	2.60	1.18	2.41	2.57	0.38	1.13

GRAIN MEAN DM% 81.4

PLOT AREA HARVESTED (OLDRESD NONE) 0.00144

PLOT AREA HARVESTED (REMAINDER) 0.00127