

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1982

[Full Table of Content](#)



## 82/R/AG/6 Agdell - W. Beans, W. Wheat

### Rothamsted Research

Rothamsted Research (1983) *82/R/AG/6 Agdell - W. Beans, W. Wheat* ; Yields Of The Field Experiments 1982, pp 26 - 30 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/AG/6

AGDELL

Object: To study, by crop yields and soil analyses, the residual values of phosphate and potash applied in the period 1848-1951 and further dressings since 1964.

The 13th year of revised scheme, w. beans, w. wheat.

For previous years see 'Details' 1967 and 1973, and 74-81/R/AG/6.

Treatments: All combinations of:-

Whole plots

1. OLDRESD      Fertilizers and organic manures applied to roots every fourth year, in the period 1848-1948:

NONE	None
PKNAMG	P K Na Mg
NPKNAMGC	N P K Na Mg C

N: 48 kg N as sulphate of ammonia

P: 41 kg P as superphosphate

K: 224 kg K as sulphate of potash

Na: 16 kg Na as sulphate of soda

Mg: 11 kg Mg as sulphate of magnesia

C: Castor meal at 2240 kg supplying about 112 kg N

2. RN CROP      Rotation 1848-1951 and crop in 1982:

F/WHEAT	With fallow: Roots (turnips or swedes), s. barley, fallow, w. wheat 1848-1951. W. wheat (after w.beans 1981)
---------	--

L/BEANS	With legume: Roots, s. barley, legume (clover or beans), w. wheat 1848-1951. W. beans (after w. wheat 1981)
---------	---

Half plots

3. 1964RESD      Residues of 1964 treatments:

P
K

Quarter plots

4. PREVCROP      Previous cropping 1958-69 on P-test half plots, 1958-70 on K-test half plots:

ARABLE	Arable or fallow
GRASS	Grass

82/R/AG/6

Sixteenth plots

5. P <sub>2</sub> O <sub>5</sub> 64	K <sub>2</sub> O 64	Rates of 1964 treatments (kg):
		P <sub>2</sub> O <sub>5</sub> to P-test      K <sub>2</sub> O to K-test
		half plots                  half plots
0	0	
500	315	
1000	630	
2000	1260	

Thirty second plots

6.		On P-test half plots: To RN CROP F/WHEAT. Residues of P <sub>2</sub> O <sub>5</sub> applied 1970-72 (kg) and in 1979 and 1981 (kg):
	P <sub>2</sub> O <sub>5</sub> 721	
	(0)0	None
	(375)300	375 total in 1970-72, 150 in 1979, 150 in 1981
		To RN CROP L/BEANS. Residues of P <sub>2</sub> O <sub>5</sub> applied 1970-72 (kg) and fresh dressings in 1980 and 1982 (kg):
	P <sub>2</sub> O <sub>5</sub> 722	
	(0)0	None
	(375)300	375 total in 1970-72, 150 in 1980, 150 in 1982
		On K-test half plots: To RN CROP F/WHEAT. Residues of K <sub>2</sub> O applied 1973-76 (kg) and in 1979 and 1981 (kg):
	K <sub>2</sub> O 761	
	(0)0	None
	(870)600	870 total in 1973-76, 300 in 1979, 300 in 1981
		To RN CROP L/BEANS. Residues of K <sub>2</sub> O applied 1973-76 (kg) and fresh dressings in 1980 and 1982 (kg):
	K <sub>2</sub> O 762	
	(0)0	None
	(870)600	870 total in 1973-76, 300 in 1980, 300 in 1982

Standard applications:

Both crops: Weedkillers: Glyphosate at 1.4 kg in 250 l. Paraquat at 0.3 kg ion and diquat at 0.3 kg ion in 250 l.  
W. beans: Weedkillers: Propyzamide at 0.85 kg in 250 l. Diquat at 0.42 kg ion (applied with 'Agral', a wetting agent) in 250 l.  
Nematicide: Aldicarb at 10 kg.

82/R/AG/6

W. wheat: Manures: 'Nitro-Chalk' at 130 kg and 770 kg. Weedkillers: Chlortoluron at 5.6 l in 250 l. Mecoprop, bromoxynil and ioxynil (as 'Brittox' at 3.5 l) in 250 l. Glyphosate at 1.4 kg in 250 l. Fungicide: Propiconazole at 0.12 kg in 250 l and on a second occasion at 0.25 kg in 250 l with the insecticide. Insecticide: Pirimicarb at 0.14 kg.

Seed: W. beans: Throws MS, sown at 300 kg.  
W. wheat: Avalon, sown at 200 kg.

Cultivations, etc.:-

Both crops: Glyphosate applied: 16 Sept, 1981. Paraquat and diquat applied: 7 Oct.  
W. beans: Aldicarb applied, seed direct drilled: 29 Oct, 1981.  
Propyzamide applied: 30 Oct. Diquat applied: 25 Aug, 1982. Combine harvested: 2 Sept.  
W. wheat: Seed direct drilled: 13 Oct, 1981. Chlortoluron applied: 16 Oct. N applied: 25 Feb, 1982 and 23 Apr. Mecoprop, bromoxynil and ioxynil applied: 19 Apr. Propiconazole applied: 26 May.  
Propiconazole with insecticide applied: 15 June. Glyphosate applied: 4 Aug. Combine harvested: 21 Aug.

82/R/AG/6

WHEAT P PLOTS

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PREVCROP	OLDRESD	NONE	(375)300	PKNAMG (0)0	(375)300	NPKNAMGC (0)0	(375)300
	P205 722 P205 64	(0)0					
ARABLE	0	6.51	7.73	7.06	6.26	3.88	3.80
	500	7.57	8.16	7.98	8.57	5.58	5.48
	1000	7.97	7.79	6.04	7.07	5.97	4.31
	2000	7.73	8.22	8.26	8.66	6.43	7.68
	0	5.27	7.44	4.65	7.37	7.06	7.73
	500	6.78	6.03	7.22	8.06	6.52	7.16
	1000	6.93	8.32	7.66	7.37	5.37	6.90
	2000	6.90	7.44	8.42	8.76	5.13	5.92

GRAIN MEAN DM% 83.6

WHEAT K PLOTS

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PREVCROP	OLDRESD	NONE	(870)600	PKNAMG (0)0	(870)600	NPKNAMGC (0)0	(870)600
	K20 762 K20 64	(0)0					
ARABLE	0	7.35	8.36	8.41	8.08	7.55	7.89
	315	7.28	7.95	8.68	8.75	8.16	8.20
	630	7.99	8.56	6.40	7.16	7.87	7.68
	1260	7.85	8.14	7.15	6.87	7.75	7.37
	0	4.73	6.81	7.73	8.43	6.97	7.98
	315	6.78	7.74	8.09	8.11	7.48	8.62
	630	7.70	8.23	7.98	8.34	6.84	7.77
	1260	7.50	8.34	8.24	8.54	6.68	7.33

GRAIN MEAN DM% 83.7

PLOT AREA HARVESTED (OLDRESD NONE) 0.00121

PLOT AREA HARVESTED (REMAINDER) 0.00136

82/R/AG/6

BEANS P PLOTS

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	OLDRESD P205 721	NONE (0)0	(375)300	PKNAMG (0)0	(375)300	NPKNAMGC (0)0	(375)300
PREVCROP	P205 64						
ARABLE	0	0.92	2.73	1.78	0.76	1.53	1.83
	500	1.98	2.93	2.37	1.95	0.47	0.54
	1000	2.48	2.99	0.45	0.99	1.61	1.53
	2000	1.40	3.57	1.60	2.57	0.85	1.00
GRASS	0	2.36	1.36	0.69	0.61	1.89	2.63
	500	1.87	0.85	0.11	0.53	1.88	1.29
	1000	1.61	0.85	0.46	0.61	2.10	1.29
	2000	2.62	1.11	0.76	0.39	2.82	2.94

GRAIN MEAN DM% 80.9

BEANS K PLOTS

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	OLDRESD K20 761	NONE (0)0	(870)600	PKNAMG (0)0	(870)600	NPKNAMGC (0)0	(870)600
PREVCROP	K20 64						
ARABLE	0	2.01	3.07	2.50	2.52	1.22	1.37
	315	3.36	3.69	2.56	3.11	0.70	0.46
	630	2.76	2.90	2.29	2.17	1.91	1.59
	1260	3.32	2.82	2.67	3.34	1.08	0.77
GRASS	0	2.61	2.11	1.15	1.22	1.04	2.44
	315	2.84	2.36	1.12	2.22	1.05	1.86
	630	2.87	1.61	2.06	0.76	0.38	0.76
	1260	2.60	1.18	2.41	2.57	0.38	1.13

GRAIN MEAN DM% 81.4

PLOT AREA HARVESTED (OLDRESD NONE) 0.00144

PLOT AREA HARVESTED (REMAINDER) 0.00127