Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1982



Full Table of Content

82/R/B/8 Plot Sizes and Mildew Spread - Spring Barley

Rothamsted Research

Rothamsted Research (1983) 82/R/B/8 Plot Sizes and Mildew Spread - Spring Barley; Yields Of The Field Experiments 1982, pp 272 - 273 - DOI: https://doi.org/10.23637/ERADOC-1-33

82/R/B/8

SPRING BARLEY

PLOT SIZES AND MILDEW SPREAD

Object: To study the effects of plot size on the incidence of mildew (Erysiphe graminis) and on the yield of neighbouring plots - Great Knott II.

Sponsor: J.F. Jenkyn.

Design: A serially balanced sequence of 4 'blocks' of 3 plots with separating and flanking plots.

Whole plot dimensions: Narrow plots: 3.0 x 12.0. Wide plots: 9.0 x 12.0.

Treatments:

TREATMNT	Plot width (all 12m long) and fungicide treatment:
3M NONE	3m, no fungicide
3M TRID	3m, tridemorph spray at 0.53 kg on 21 May
9M NONE	9m, no fungicide

- NOTES: (1) The above plots were each separated by 3m wide plots sprayed with tridemorph. Three extra plots, also treated with tridemorph, were included at each end of the experiment.
 - (2) The effects of treatments to neighbouring plots (left LHN, right - RHN) were estimated. In this experiment 'left' was East, 'right' was West.
- Basal applications: Manures: (0:14:28) at 900 kg, 'Nitro-Chalk' at 480 kg. Weedkillers: Dicamba, mecoprop and MCPA (as 'Poly-Farmon' at 5.0 1) in 250 l. Paraquat and diquat (as 'Cleansweep' at 2.0 1) in 250 l.

Seed: Georgie, sown at 160 kg.

Cultivations, etc.:- Paraquat and diquat applied: 22 Oct, 1981. PK applied: 25 Nov. Ploughed: 10 Dec. Spring-tine cultivated, N applied: 30 Mar, 1982. Rotary harrowed, seed sown: 31 Mar. Dicamba, mecoprop and MCPA applied: 18 May. Combine harvested: 17 Aug. Previous crops: S. beans 1980, w. barley 1981.

NOTE: Mildew was assessed in early June and early July on all plots.

82/R/B/8

**** TABLES OF MEANS ****

	TREA	TMMT	314	NONE 5.72	3M	TRID 6.62	9M	NONE 5.39
	TRE	LHN	3M	NONE	3M	TRID	9M	NONE
	3M	NONE				5.64		5.80
		TRID		6.61 5.41		5.38		6.64
	TOF	RHN	3M	NONE	3M	TRID	9M	NONE
TREATMNT					c 77		F 67	
		NONE		c 70		5.77		5.67
		TRID		6.70				6.55
	9M	NONE		5.53		5.26		
	GRAND	MEAN		5.91				

GRAIN MEAN DM% 84.0

PLOT AREA HARVESTED 0.00245