

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1982

[Full Table of Content](#)



## 82/R/O/1 Aldicarb and Stem Nematode - Spring Oats

### Rothamsted Research

Rothamsted Research (1983) *82/R/O/1 Aldicarb and Stem Nematode - Spring Oats* ; Yields Of The Field Experiments 1982, pp 276 - 277 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/0/1

SPRING OATS

ALDICARB AND STEM NEMATODE

Object: To study the effects of a range of rates and methods of applying aldicarb on the control of stem nematode (*Ditylenchus dipsaci*) and on the yield of s. oats - Fosters 0 and E VI.

Sponsor: A.G. Whitehead.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 2.29 x 7.32.

Treatments:

TREATMNT	Aldicarb rates and row spacings:
	Sown in rows 12.7 cm (5 inches) apart:
0 CR	No aldicarb
2 CR	Aldicarb at 1.25 kg per ha, applied equally to every row
4 CR	Aldicarb at 2.5 kg per ha, applied equally to every row
1 CR ALT	Aldicarb at 0.6 kg per ha, applied to alternate rows only
2 CR ALT	Aldicarb at 1.25 kg per ha, applied to alternate rows only
4 CR ALT	Aldicarb at 2.5 kg per ha, applied to alternate rows only
	Sown in 7.6 cm (3 inches) wide bands, centres of bands 25.4 cm (10 inches) apart:
0 WB	No aldicarb
1 WB	Aldicarb at 0.6 kg per ha, applied equally to all bands
2 WB	Aldicarb at 1.25 kg per ha, applied equally to all bands
4 WB	Aldicarb at 2.5 kg per ha, applied equally to all bands

Basal applications: Manures: (20:10:10) at 360 kg. 'Nitro-Chalk' at 140 kg. Weedkillers: Mecoprop, bromoxynil and ioxynil (as 'Brittox' at 3.5 l) in 220 l.

Seed: Maris Osprey, sown at 180 kg.

Cultivations, etc.: - Ploughed: 15 Oct, 1981. NPK applied: 23 Mar, 1982. Treatments applied, rotary harrowed, seed sown: 25 Mar. Weedkillers applied: 13 May. N applied by hand: 28 May. Combine harvested: 10 Aug. Previous crops: Mixed crops 1980, w. oats 1981.

82/R/0/1

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT	
0 CR	3.47
2 CR	3.57
4 CR	3.59
1 CR ALT	3.34
2 CR ALT	3.44
4 CR ALT	3.67
0 WB	2.95
1 WB	3.13
2 WB	3.27
4 WB	3.17
MEAN	3.36

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT
-----	-----
SED	0.139

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	18	0.171	5.1

GRAIN MEAN DM% 88.2

PLOT AREA HARVESTED 0.00093 (ROWS 12.7CM SPACING)  
0.0111 (ROWS 7.6CM SPACING)