

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1982

[Full Table of Content](#)



### 82/R/BE/1 Effects of Pests and Pathogens - Winter Beans

#### Rothamsted Research

Rothamsted Research (1983) *82/R/BE/1 Effects of Pests and Pathogens - Winter Beans* ; Yields Of The Field Experiments 1982, pp 278 - 279 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/BE/1

WINTER BEANS

EFFECTS OF PESTS AND PATHOGENS

Object: To assess the benefits from three amounts of pest and disease control on beans - Gt. Knott III.

Sponsors: J. McEwen, A. Bainbridge, R. Bardner, A.J. Cockbain, J.M. Day, K.E. Fletcher, D.C. Griffiths, D.H. Lapwood, R.M. Webb, T.D. Williams, D.P. Yeoman.

Design: 6 randomised blocks of 3 plots.

Whole plot dimensions: 5.33 x 15.0.

Treatments:

PATHCONT	Pest and pathogen control (in addition to basals):
STANDARD	None
ENHANCED	Seed dressed with benomyl and thiram (1.2 g of each per kg of seed)
FULL	Phorate at 2.2 kg as granules to foliage on 2 Apr, 1982 Seed dressed with benomyl and thiram (1.2 g of each per kg of seed) Aldicarb at 10 kg on 23 Sept, 1981 Benomyl at 0.56 kg and fosetyl-Al at 2.2 kg on 8 Feb Carbofuran at 2.24 kg on 2 Apr Benomyl at 0.56 kg on 14 Apr Benomyl at 0.56 kg on 5 May Propiconazole at 0.13 kg on 15 June

NOTES: (1) Treatment sprays were applied in 340 l.  
(2) Sides of plots were separated by strips of w. beans 5.33 m wide plus 0.66 m fallow paths, ends of plots were separated by strips of w. beans 9.2 m wide plus 0.9 m fallow paths. The separating crops received basal applications as for the plots and in addition received benomyl at 0.56 kg on 8 Feb and 14 Apr.

Basal applications: Weedkillers: Paraquat at 0.56 kg ion in 220 l.  
Propyzamide at 0.85 kg in 250 l. Fungicide: Benomyl at 0.55 kg in 250 l on two occasions.

Seed: Throws MS, sown at 250 kg.

Cultivations, etc.: - Heavy spring-tine cultivated twice: 9 Sept, 1981 and once: 11 Sept. Paraquat applied: 22 Sept. Spring-tine cultivated, seedbed treatments applied, rotary harrowed, seed sown: 23 Sept. Propyzamide applied: 29 Sept. Basal benomyl applied twice: 19 May, 1982, 7 June. Combine harvested: 12 Aug. Previous crops: W. wheat 1980, s. barley 1981.

NOTE: Plant counts were made after establishment and components of yield were measured at maturity. Total above-ground dry matter and N content were measured in July. Migratory nematodes, root and foliar fungi, aphids, weevils, midges and viruses were counted at intervals during the season. N content of grain was measured.

82/R/BE/1

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PATHCONT	STANDARD	ENHANCED	FULL	MEAN
	2.97	2.99	3.04	3.00

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	PATHCONT
-----	-----
SED	0.155

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.268	8.9

GRAIN MEAN DM% 85.1

PLOT AREA HARVESTED 0.00320