

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1982

[Full Table of Content](#)



### 82/R/BE/2 Control of Chocolate Spot - Winter Beans

#### Rothamsted Research

Rothamsted Research (1983) *82/R/BE/2 Control of Chocolate Spot - Winter Beans* ; Yields Of The Field Experiments 1982, pp 280 - 281 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/BE/2

WINTER BEANS

CONTROL OF CHOCOLATE SPOT

Object: To study the effects of times and frequencies of application of two fungicides on the control of chocolate spot and on the yield of w. beans - Gt. Knott III.

Sponsors: A. Bainbridge, G.R. Cayley.

Design: 3 randomised blocks of 16 plots.

Whole plot dimensions: 5.33 x 10.0.

Treatments: All combinations of:-

1. FUNGCIDE Fungicides:

BENOMYL	Benomyl at 0.56 kg
PROCHLOR	Prochloraz at 0.50 kg

2. FUNGTIME Times and frequencies of applying fungicides:

1+5	Twice, on 25 Jan, 1982 and 11 May
5+7	Twice, on 11 May and 9 June
2	Once, on 19 Mar
3	Once, on 22 Apr
4	Once, on 29 Apr
6	Once, on 26 May
7	Once, on 9 June

plus one extra treatment:

EXTRA

NONE None (duplicated)

Basal applications: Weedkillers: Propyzamide at 0.85 kg in 250 l.  
Paraquat at 0.56 kg ion in 220 l.

Seed: Throws MS, sown at 250 kg.

Cultivations, etc.:- Deep spring-tine cultivated twice: 9 Sept, 1981 and a third time: 11 Sept. Paraquat applied: 22 Sept. Spring-tine cultivated, seed sown: 23 Sept. Propyzamide applied: 29 Sept. Combine harvested: 12 Aug, 1982. Previous crops: W. barley 1980, s. barley 1981.

NOTE: Emergence counts were made in autumn and components of yield measured at maturity. Chocolate spot was assessed monthly throughout the growing season and leaf drop counts were made in mid-June.

82/B/BE/2

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

FUNGCIDE FUNGTIME	BENOMYL	PROCHLOR	MEAN
1+5	2.73	2.49	2.61
5+7	2.60	2.38	2.49
2	2.36	2.36	2.36
3	2.44	2.49	2.47
4	2.58	2.41	2.50
6	2.94	2.44	2.69
7	2.45	2.48	2.47
MEAN	2.59	2.44	2.51

EXTRA NONE 2.45

GRAND MEAN 2.50

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	FUNGTIME	FUNGCIDE	FUNGTIME FUNGCIDE
-----	-----	-----	-----
SED	0.078	0.042	0.111

SED FOR COMPARING EXTRA NONE WITH ANY ITEM IN FUNGTIME.FUNGCIDE TABLE IS 0.096

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	31	0.135	5.4

GRAIN MEAN DM% 86.0

PLOT AREA HARVESTED 0.00320