

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/BE/7 Effects of Pests and Pathogens - Spring Beans

Rothamsted Research

Rothamsted Research (1983) *82/R/BE/7 Effects of Pests and Pathogens - Spring Beans* ; Yields Of The Field Experiments 1982, pp 290 - 291 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/BE/7

SPRING BEANS

EFFECTS OF PESTS AND PATHOGENS

Object: To assess the benefits from three amounts of pest and disease control on irrigated and unirrigated s. beans - W. Barnfield I.

Sponsors: J. McEwen, R. Bardner, A.J. Cockbain, J.M. Day, K.E. Fletcher, D.H. Lapwood, B.J. Legg, R.M. Webb, T.D. Williams, J.F. Witty, D.P. Yeoman.

Design: 4 randomised blocks of 2 plots split into 3.

Whole plot dimensions: 4.27 x 13.7.

Treatments: All combinations of:-

Whole plots

1. IRRIGATN Irrigation:
- | | |
|------|--------------------|
| NONE | None |
| FULL | Full (total 25 mm) |

Sub plots

2. PATHCONT Pest and pathogen control:
- | | |
|----------|---|
| STANDARD | None |
| ENHANCED | Phorate at 2.2 kg, combine drilled
Pirimicarb at 0.14 kg on 15 June, 1982
Benomyl at 0.50 kg on 2 July |
| FULL | Aldicarb at 10 kg on 24 Mar
Phorate at 2.2 kg, combine drilled
Fosetyl-Al at 2.2 kg on 1 June
Pirimicarb at 0.14 kg on 15 June
Benomyl at 0.50 kg on 2 July
Benomyl at 0.56 kg on 30 July
Propiconazole at 0.13 kg on 30 July and on 13 Aug |

Basal applications: Weedkillers: Trietazine at 1.0 kg with simazine at 0.14 kg in 250 l.

Seed: Minden, sown at 230 kg.

Cultivations, etc.:- Disced: 24 Sept, 1981. Ploughed: 29 Jan, 1982. Spring-tine cultivated twice: 23 Mar, 24 Mar. Rotary harrowed: 24 Mar. Seed sown: 25 Mar. Weedkillers applied: 27 Mar. Combine harvested: 2 Sept. Previous crops: S. barley 1980 and 1981.

NOTE: Plant counts were made after establishment and components of yield were measured at maturity. Total above-ground dry matter and N content were measured in August. Migratory nematodes, root and foliar fungi, aphids, weevils and viruses were counted at intervals during the season. N content of grain was measured.

82/R/BE/7

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PATHCONT IRRIGATN	STANDARD	ENHANCED	FULL	MEAN
NONE	3.52	3.94	4.38	3.95
FULL	3.80	4.09	4.41	4.10
MEAN	3.66	4.01	4.40	4.02

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PATHCONT	IRRIGATN* PATHCONT

SED	0.085	0.120

* WITHIN THE SAME LEVEL OF IRRIGATN ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	12	0.170	4.2

GRAIN MEAN DM% 81.7

SUB PLOT AREA HARVESTED 0.00293