

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/BE/8 Varieties and Blrv - Spring Beans

Rothamsted Research

Rothamsted Research (1983) *82/R/BE/8 Varieties and Blrv - Spring Beans* ; Yields Of The Field Experiments 1982, pp 292 - 293 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/BE/8

SPRING BEANS

VARIETIES AND BLRV

Object: To study the effects of full pest and pathogen control on four varieties differing in susceptibility to bean leaf roll virus (BLRV) - Claycroft.

Sponsor: A.J. Cockbain.

Design: 3 randomised blocks of 8 plots.

Whole plot dimensions: 4.27 x 9.14.

Treatments: All combinations of:-

1. VARIETY Varieties:
 BEAD Maris Bead (field bean, resistant to BLRV)
 MINDEN Minden (field bean, sensitive to BLRV)
 T WHITE Threefold White (broad bean, sensitive to BLRV)
 WEIRBOON Weirboon (broad bean, tolerant to BLRV)
2. PATHCONT Pest and pathogen control:
 STANDARD None
 FULL Phorate, pirimicarb and benomy1

NOTE: Rates of materials used in PATHCONT FULL were
 Phorate at 4.5 kg and rotary harrowed in to the seedbed on
 25 Mar, 1982
 Pirimicarb at 0.14 kg in 340 l on 15 June
 Benomy1 at 0.5 kg in 340 l on 2 July

Basal applications: Manures: Chalk at 5.0 t. Weedkillers: Glyphosate at 1.4 kg in 250 l. Trietazine at 1.0 kg with simazine at 0.14 kg in 250 l.

Cultivations, etc.:- Glyphosate applied: 4 Nov, 1981. Chalk applied: 10 Nov. Ploughed: 27 Nov. Spring-tine cultivated: 24 Mar, 1982. Seed sown: 26 Mar. Trietazine with simazine applied: 1 Apr. Combine harvested: 2 Sept. Previous crops: W. wheat 1980 and 1981.

NOTE: Plant counts were made at emergence. Pest and disease incidence and growth parameters were assessed throughout the season.

82/R/BE/8

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

VARIETY	BEAD	MINDEN	T WHITE	WEIRBOON	MEAN
PATHCONT					
STANDARD	4.04	3.89	3.77	4.31	4.00
FULL	4.49	4.52	4.27	4.22	4.38
MEAN	4.26	4.20	4.02	4.27	4.19

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	PATHCONT	VARIETY PATHCONT

SED	0.187	0.132	0.264

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	14	0.323	7.7

GRAIN MEAN DM% 78.8

PLOT AREA HARVESTED 0.00293