

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/BE/12 Control of Rust - Spring Beans

Rothamsted Research

Rothamsted Research (1983) *82/R/BE/12 Control of Rust - Spring Beans* ; Yields Of The Field Experiments 1982, pp 298 - 299 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/BE/12

SPRING BEANS

CONTROL OF RUST

Object: To study the effects of fungicides on the control of rust (*Uromyces fabae*) and on the yield of spring beans - Long Hoos VI/VII 5.

Sponsors: D.H. Lapwood, J. McEwen, D.P. Yeoman.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 2.03 x 3.05.

Treatments: All combinations of:-

1. C S FUNG Fungicide to control chocolate spot but not rust:
 NONE None
 BENOMYL Benomyl at 1.0 kg in 340 l on 2 July, 1982
2. RUSTFUNG Fungicides to control rust:
 MAN+MANC Maneb at 0.8 kg + mancozeb at 0.8 kg in 340 l
 PROPICON Propiconazole at 0.12 kg in 340 l
3. RFNGTIME Times of applying fungicides to control rust:
 TWICE Twice on 9 July and 13 Aug
 THRICE Thrice, on 9 July, 23 July and 13 Aug

plus one extra treatment:

EXTRA

NONE None (duplicated)

Basal applications: Manures: Chalk at 2.9 t. Muriate of potash at 520 kg.
Weedkillers: Trietazine at 1.0 kg with simazine at 0.14 kg in 340 l.
Insecticide: Permethrin applied twice at 0.06 kg in 340 l.

Seed: Minden, sown at 270 kg.

Cultivations, etc.: - Muriate of potash applied: 12 Nov, 1981. Chalk applied: 27 Nov. Ploughed: 29 Jan, 1982. Spring-tine cultivated: 26 March. Spring-tine cultivated, power harrowed, seed sown: 29 Mar. Weedkillers applied: 14 Apr. Insecticide applied: 11 May, 26 May. Harvested by hand: 9 Sept. Previous crops: Potatoes 1980, fallow 1981.

NOTE: The incidence of chocolate spot and rust was assessed from early July until maturity. Components of yield were measured at maturity.

82/R/BE/12
GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

RUSTFUNG	MAN+MANC	PROPICON	MEAN
C S FUNG			
NONE	5.28	5.18	5.23
BENOMYL	5.65	5.60	5.62
MEAN	5.46	5.39	5.43

RFNGTIME	TWICE	THRICE	MEAN
C S FUNG			
NONE	5.06	5.40	5.23
BENOMYL	5.47	5.77	5.62
MEAN	5.27	5.59	5.43

RFNGTIME	TWICE	THRICE	MEAN
RUSTFUNG			
MAN+MANC	5.36	5.57	5.46
PROPICON	5.17	5.61	5.39
MEAN	5.27	5.59	5.43

C S FUNG	RFNGTIME	TWICE	THRICE
NONE	RUSTFUNG		
	MAN+MANC	5.05	5.50
	PROPICON	5.06	5.30
BENOMYL	MAN+MANC	5.67	5.63
	PROPICON	5.28	5.92

EXTRA NONE 4.51

GRAND MEAN 5.24

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	C S FUNG	RUSTFUNG	RFNGTIME	C S FUNG RUSTFUNG
SED	0.138	0.138	0.138	0.195

TABLE	C S FUNG RFNGTIME	RUSTFUNG RFNGTIME	C S FUNG RUSTFUNG RFNGTIME
SED	0.195	0.195	0.276

SED FOR COMPARING EXTRA NONE WITH ANY ITEM IN
C S FUNG.RUSTFUNG.RFNGTIME TABLE IS 0.239

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	19	0.338	6.5
GRAIN MEAN DM%	87.0	PLOT AREA HARVESTED	0.00015